

Singlet-triplet Splitting and Stability of Saturated Acyclic and Cyclic Carbenes: DFT Calculations

ESMAIL VESSALLY

Payame Noor University (PNU), Zanjan, Iran.

(Received 14th May 2008, accepted in revised form 12th September 2008)

Summary: DFT calculations were carried out on the singlet-triplet splitting for $C_nH_{2n+2}C:$ ($n = 0-7$) and $C_nH_{2n}C:$ ($n = 2-7$) at B3LYP/6-311++G (3df,2p) level of theory. For the estimation of stabilization energies (SE), the isodesmic reactions were applied. The results of the isodesmic reaction show a high stability for singlet state relative to the corresponding triplet state by substitution of methyl or methylene groups at the both sides of methylenic center in $C_nH_{2n+2}C:$. Substitution of methylene groups at the both sides of cyclopropenylidene ($C_nH_{2n}C:$) leads to reduce the instability of triplet state; decreasing the singlet-triplet splitting.


Introduction

Highly reactive carbenes intermediates have attracted much attention in organic chemistry [1-5]. The reactivity of carbenes can be understood in terms of their electronic structure; triplet state ($\sigma\pi$ configuration) and singlet states (σ^2 , π^2 and $\sigma\pi$ configurations) (Scheme-1). The trends in the reactivity of carbenes can be recognized through determination of their singlet-triplet splitting. For both the triplet and the singlet states, the π -orbital is essentially a pure p -like orbital while the σ -orbital is a hybrid of s - and p -like atomic orbital. Thus, σ -orbital with increased p -character favor a triplet ground state while the σ -orbital with increased s -character favor singlet state. The hybridization of the σ -orbital is in turn determined by the following factors [6]. First, the bulky substituents favor a large bond angle. This leads to decreasing s -character in the nonbonding σ -orbital, thus favoring the triplet state. Secondly, the substituents with high electronegativity respect to the carbon form a polar bond. The polar bond increases the p -orbital contributions, which in turn decreases the p -character in the

nonbonding carbon σ -orbital. As a result, the σ -orbital increases the s character and stabilized the singlet state. In contrast, the electropositive substituents favor the triplet state. Thirdly, hyperconjugation also increase the singlet-triplet splitting with stabilizing the singlet state while destabilizing the triplet state. Valence electron pairs on the substituent may donate electron density to the nonbonding σ -orbital of the central carbon. Thus, the singlet state should be stabilized by substituents that are electronegative and/or have electron pairs that can be donated *via* hyperconjugation to the empty σ -orbital. Our goal in the present work is to obtain quantitative information on the magnitude of the relative stability and singlet-triplet splitting of the simple saturated acyclic $C_nH_{2n+2}C:$ and cyclic $C_nH_{2n}C:$ carbenes.

Results and Discussion

As the continuation of our previous works, our goal of investigation is to obtain quantitative



Scheme-1: Electronic structures of triplet ($\sigma\pi$ configuration) state and singlet states (σ^2 , π^2 and $\sigma\pi$ configurations).

methylene groups at the both sides of cyclopropenylidene ($C_nH_{2n}C$) leads to reduce the instability of triplet state; decreasing the singlet-triplet splitting.

Experimental

Full geometry optimizations were carried out by B3LYP [8, 9] methods using 6-311++G(3df,2p) basis set of the Gaussian 98 system of programs [10]. The geometry will be adjusted until a stationary point on the potential surface was found. Here, the Berny algorithm was employed for all minimizations using redundant internal coordinates [11]. For minimum state structures, only real frequency values were considered.

Conclusion

For $C_nH_{2n+2}C$, the methylene ($:CH_2$) was a ground state triplet while substitution of one or two methyl groups at methylenic center leads to reduce the singlet-triplet splitting with the singlet ground state. Substitution of three and four methyl groups on methylenic center leads to raising the singlet-triplet splitting. The results of the isodesmic reaction show more stability of the singlet state relative to triplet state by substitution of more methyl groups at methylenic center for $C_nH_{2n+2}C$. For $C_nH_{2n}C$, the cyclopropenylidene have a ground state singlet. Substitution of a methylene group on cyclopropenylidene (C_3H_6C) leads to reduce the singlet-

triplet splitting. Substitution of methylene groups at the both sides of cyclopropenylidene ($C_nH_{2n}C$) leads to reduce the instability of triplet state; decreasing the singlet-triplet splitting.

References

1. P. P. Gaspar, M. Xiao, D. H. Pae, D. J. Berger, T. Haile, T. Chen, D. Lei, W. R. Winchester and P. Jiang, *J. Organomet. Chem.*, **646**, 68 (2002).
2. H. Wang and K. Brezinsky, *J. Phys. Chem. A*, **102**, 1530 (1998).
3. M. Z. Kassae, B. N. Haerizade and S. Arshadi, *J. Mol. Struct. (Theochem.)*, **639**, 187 (2003).
4. K. Klinkhammer, *Polyhedron*, **21**, 587 (2002).
5. M. Denk, R. Lennon, R. Hayashi, R. West, A. V. Belyakov, H. P. Verne, A. Haaland, M. Wagner and N. Metzler, *J. Am. Chem. Soc.*, **116**, 2691 (1994).
6. E. A. Carter and W. A. Goddard III, *J. Chem. Phys.*, **88**, 1752 (1988).
7. C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B*, **37**, 785 (1988).
8. A. D. Becke, *J. Chem. Phys.*, **98**, 5648 (1993).
9. GAUSSIAN 98, Revision A.6, M. J. Frisch, et al., Gaussian Inc., Pittsburgh, PA, (1998).
10. C. Peng, P. Y. Ayala, H. B. Schlegel and M. J. Frisch, *J. Comput. Chem.*, **17**, 49 (1996).
11. M. K. Cyranski, P. V. R. Schleyer, T. M. Krygowski, H. Jiao and G. Hohlneicher, *Tetrahedron*, **59**, 1657 (2003).