

Interaction of Nitroglycerin with Bovine Serum Albumin and the Influence of Metal Ions on the Binding

Chengman Bao, Jialian Wang, Xuehong Tong, Chunli Zhang, Xinhui Tang*

¹College of Ocean and Biological Engineering, Yancheng Teachers University, Yancheng 224051, PR China. bbccmm_hi@126.com*

(Received on 20th November 2018, accepted in revised form 27th June 2019)

Summary: The effect of Cu²⁺, Ca²⁺, Mg²⁺ and Zn²⁺ on the interaction between nitroglycerin and bovine serum albumin was investigated. The bimolecular quenching rate constant, the Stern-Volmer quenching constant, the binding constants and the number of binding sites were calculated in the absence and presence of Cu²⁺, Ca²⁺, Mg²⁺ and Zn²⁺. The quenching constants of nitroglycerin to bovine serum albumin were increased in the presence of metal ions. Static quenching mechanism was also confirmed. The binding constants of nitroglycerin to bovine serum albumin were influenced by different metal ions. The enthalpy change, free energy change, entropy change and the distance between the donor and the acceptor at different temperatures were calculated. The results indicated that energy transfer from bovine serum albumin to nitroglycerin occurs with high probability.

Keywords: Nitroglycerin; Bovine serum albumin; Metal ions; Spectroscopy.

Introduction

An important property of serum albumins is transporting compounds like drugs and pharmaceuticals [1-2].

The transport mechanism of drugs in plasma could provide important information for drug development. Bovine serum albumin (BSA) always as an important protein to study the drugs in human body for its approximately 76% sequence homologous with human serum albumin (HSA) [3-4]. In human organism, some metal ions are important to proteins. For albumin, about 50% of Ca²⁺ ions, 10% of Cu²⁺ ions, 10% of Zn²⁺ ions are carried in the serum [5]. The binding sites or conformational changes between drug and serum albumin always influenced by metal ions which can react with serum albumins [6-10]. Some researches have been showed that metal ions affected significantly the drug-serum albumin interactions [11].

Nitroglycerin (NTG) is a medication used to treat of various diseases such as angina and tendinopathies [12-18]. Many methods such as fluorescence spectroscopy and ultraviolet spectroscopy have been widely used to study the binding of drugs with protein [19-24]. Herein, the interaction between nitroglycerin with bovine serum albumin (BSA) was studied using the fluorescence spectroscopy and ultraviolet spectroscopy. Four metal ions, Cu²⁺, Ca²⁺, Mg²⁺ and Zn²⁺ were also selected to gain insight the influence of metal ions on the interaction.

Experimental

Materials and apparatus

BSA and diluted nitroglycerin were purchased from Sigma-Aldrich. Zn²⁺, Cu²⁺, Ca²⁺, Mg²⁺ are come from their chloride. The purity of the chemicals (EDTA and Tris-HCl) were 99%. The effect of Cl⁻ can be ignored with Tris-HCl [25].

RF-5301PC (SHIMADZU) spectrofluorometer which equipped with 1.0 cm quartz cells was used to recorded the fluorescence spectra. UV-vis spectrophotometer (UV-1800, SHIMADZU) equipped with 1.0 cm quartz cells was used to record the UV spectra.

Spectra measurement of BSA and NTG in the absence and presence of metal ions

The excitation wavelength in our experiments was 285 nm and the ranges of spectra are from 300 nm to 450 nm. The excitation and emission slit widths were 3 nm. Stock solution of BSA (1×10⁻⁴ M) was dissolved in 0.05 M Tris-HCl buffer (pH 7.4, containing 0.1 M NaCl). Nitroglycerin were further diluted in Tris-HCl buffer (pH 7.4) at 25 °C. Samples containing 2 ml of the BSA solution and NTG solution were mixed. The final concentrations of NTG were from 50 μM to 200 μM. The ranges of absorption spectra of protein were of 200nm to 450 nm.

The stock solution of different metal ions

*To whom all correspondence should be addressed.

(1.0×10^{-2} M) was prepared by dissolving in Tris-HCl buffer. The concentrations of BSA which we used in our experiments were 1×10^{-6} M.

Results and Discussion

Quenching mechanism of BSA fluorescence by NTG

The effect of NTG on the BSA fluorescence spectra is shown in Fig 1. The fluorescence intensity of BSA was gradually decreased upon increasing the concentration of NTG. The decrease in fluorescence intensity indicates that there are interactions between NTG and BSA.

A decrease of the fluorescence intensity of a compound can be due to molecular rearrangements and collisional quenching [23]. The Stern-Volmer equation is used to describe fluorescence quenching [24]:

$$F_0 / F = 1 + Kq\tau_0[Q] = 1 + K_{sv}[Q] \quad (1)$$

F_0 , the relative fluorescence intensity of protein. F , the relative fluorescence intensity of donor when quencher is exist. Kq , the bimolecular quenching rate constant. K_{sv} , the Stern-Volmer quenching constant. τ_0 , the average life time of the biomolecule without quencher ($\tau_0 = 10^{-8}$ s) [25], $[Q]$, the concentration of the quencher. The curves of F_0/F versus $[Q]$ is shown in Fig 2.

Fig 2 showed a good linear relationship during the addition of NTG 100-300 μ M. The dynamic Ster-Volmer quenching constants (K_{sv}) and quenching rate constants Kq were calculated based on the fluorescence life time of biopolymers (Table-1).




Fig 1: Fluorescence spectra of BSA. (1→4) pure BSA, with NTG at 100 μ M, with NTG at 200 μ M, and with NTG at 300 μ M.

Table-1. The values of Stern-Volmer constants and binding parameters at different temperatures.

T (K)	Kq ($\times 10^{12}$ L/mol·S)	K_{sv} ($\times 10^4$ L/mol)	Ka ($\times 10^3$ L/mol)	n
288	0.138	0.138	3.8	0.8822
298	0.057	0.057	0.88	0.8229
308	0.058	0.058	69.31	1.2584




Fig 2: Stern-Volmer curves of NTG quenching BSA fluorescence at (1) 288 K, (2) 298 K, and (3) 308 K.

The maximum collisional quenching constant of different quenchers with biomolecule is $2.0 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ [26]. The bimolecular quenching rate constants (Kq) are larger than the maximum collisional quenching constant. The main mechanism of the quenching could be the static quenching. Some researches have been proved that other proteins have more distinct photoluminescence exhibiting groups [27]. Fluorescence quenching polymer (Ppy) and the flavin adenine dinucleotide (FAD) composite with Ppy effectively quenched the FAD fluorescence [28]. A fluorescence quenching polymer (Ppy) was used to increase the selectivity and sensitivity of an immunoanalytical system as fluorescence quencher [29]. These results can provide useful information for the application of fluorescence studies.

Small molecules are assumed to bind independently to a set of equivalent sites on a macromolecule [30]. The number of binding sites and the binding constant of the interaction between NTG and BSA were described with the following equation [31].

$$\log[(F_0 - F)/F] = \log Ka + n \log [Q] \quad (2)$$

n , is the number of binding sites, Ka , the binding constant. The $\log [(F_0 - F)/F]$ versus $\log [Q]$ were showed in Fig 3.

The values of binding parameters are given in Table-1. There is one binding site in NTG to BSA

for the values of n were approximately equal to 1. Large values of Ka show strong binding between NTG and BSA [32]. The results showed that sequence of the binding strength was $308 \text{ K} > 288 \text{ K} > 298 \text{ K}$.




Fig 3: Plot of $\log[(F_0 - F)/F]$ versus $\log[Q]$ for quenching of BSA by NTG at (1) 288 K, (2) 298 K, and (3) 308 K

Thermodynamic parameters

Electrostatic interactions, hydrogen bonds, hydrophobic forces, and Van der Waals always to explain the mechanisms of the combine of small molecules to macromolecules [9,33]. The thermodynamic parameters, Gibbs free energy changes (ΔG), enthalpy changes (ΔH), entropy change (ΔS) were calculated to clarify the interaction of NTG with BSA by the following equations.

$$\ln Ka = -\Delta H / RT + \Delta S / R \quad (3)$$

$$\Delta G = -RT \ln Ka \quad (4)$$

$$\Delta S = (\Delta H - \Delta G) / T \quad (5)$$

R , the gas constant. T , temperature in Kelvin scale. Ka , the equilibrium constant at temperature T . ΔH and ΔS can be obtained from the slope and intercept of the plot of $\ln Ka$ versus $1/T$. The value of ΔG can be calculated using Eq.(4). The results are shown in Table-2. For $\Delta G < 0$ means that the interaction process of NTG is spontaneous. Researches have been proved that the thermodynamic parameters associated with various interaction that may take place in

ligand-protein binding process: $\Delta S > 0$ and $\Delta H > 0$ means the main binding force was hydrophobic force, $\Delta H < 0$ and $\Delta S < 0$ means the main binding force were vander Waals' force and hydrogen bonding, $\Delta H < 0$ and $\Delta S > 0$, means the main binding force was electrostatic interactions [31]. Therefore, the positive ΔS and ΔH means the main binding force between NTG and BSA was hydrophobic force.

Table-2: Thermodynamic parameters for BSA-NTG system.

T (K)	ΔH (KJ/mol)	ΔS (J/mol)	ΔG (KJ/mol)
288			-19.733
298	104.698	424.103	-16.789
308			-28.543

Energy transfer between NTG and BSA

According to the Förster theory, the energy transfer efficiency E is defined as the following Eq.(6).

$$E = 1 - F/F_0 = R_0^6 / (R_0^6 + r^6) \tag{6}$$

r , the distance from the donor to the acceptor, R_0 , the Förster critical distance (when the efficiency of energy transfer is 50%). R_0 can be calculated using following equations.

$$R_0^6 = 8.8 \times 10^{-25} K^2 \Phi N^{-4} J \tag{7}$$

$$J = [\sum F(\lambda)\epsilon(\lambda)\lambda^4 \Delta\lambda] / [\sum F(\lambda)\Delta\lambda] \tag{8}$$

K^2 , the space factor orientation. N , the average refractive index of medium respectively ($K^2=2/3$, $N=1.336$ for BSA [32].). In Eq.(8), $F(\lambda)$ is the corrected fluorescence intensity of the donor in the wavelength range λ , $\epsilon(\lambda)$ is the molar absorptivity of the acceptor at wavelength λ . J is the spectral overlap between the emission spectrum of the donor and the absorption spectrum of the acceptor.

Fig 4 shows the spectral overlap between absorption spectrum of NTG with fluorescence spectrum of BSA in the wavelength range of 300nm to 450 nm. Table-3 shows BSA to NTG distance in the complex is less than 7 nm, indicating that energy transfer from BSA to NTG occurs with high probability.

Table-3. Parameters of E , J , R_0 and r of NTG with BSA under different temperatures.

T (K)	E	J ($\times 10^{14}$ cm ³ ·L/mol)	R_0 (nm)	r (nm)
288	0.132	6.657	3.435	4.7
298	0.064	6.88	3.454	5.387
308	0.034	6.865	3.453	6.016

The interactions between NTG and BSA in the presence of different metal ions.

Metal ions play significant role in human organism. Many researches have been studied the interactions between serum albumin and some drugs in the presence of metal ions [34-36]. The results showed that the binding constants of drugs and BSA was changed by metal ions [37].

As shown in Fig 5, when different metal ions were titrated into NTG-BSA solution, the maximum of λ_{em} and shape of the fluorescence spectra of BSA-NTG were no obvious change to those in the absence of metal ions. But the fluorescence quenching extent was larger than those without metal ions. The quenching extent of different metal ions was $Cu^{2+} > Ca^{2+} > Mg^{2+} > Zn^{2+}$. The results suggested that the binding of NTG with BSA was influenced by metal ions.

The curves of F_0/F versus $[Q]$ were shown in Fig 6. Fig 6 showed a good linear relationship for NTG with BSA in the presence of metal ions. The values of Kq for the NTG in the presence of different metal ions were calculated and were higher than the value of the maximum scatter collision quenching constant (Table-4). The results suggested that the formation of a complex between NTG and BSA, corresponding to a static quenching mechanism no matter metal ions was absent or present in the system. The quenching constants (K_{sv}) of NTG to BSA were increased in the presence of metal ions and the extent was $Ca^{2+} > Cu^{2+} > Zn^{2+} > Mg^{2+}$.

Table-4: The Stern-Volmer quenching constant, quenching rate constant, binding constants and binding sites of NTG-BSA in the presence of different metal ions at 298 K

BSA-NTG-M ²⁺	K_{sv} ($\times 10^4$ L/mol)	Kq ($\times 10^{12}$ L/mol·S)	Ka ($\times 10^3$ L/mol)	n
BSA-NTG-Mg ²⁺	0.115	0.115	5.05	0.9276
BSA-NTG-Cu ²⁺	0.1856	0.1856	25.12	1.0293
BSA-NTG-Ca ²⁺	0.1915	0.1915	0.83	0.7264
BSA-NTG-Zn ²⁺	0.1397	0.1397	0.62	0.7253




Fig 4: The spectral overlap between absorption spectrum of NTG (1) with fluorescence spectrum of BSA (2) under different temperatures.




Fig 5: The fluorescence spectra of BSA-NTG in the absence and presence of metal ions. 1→5 : NTG-BSA, NTG-BSA-Zn²⁺, NTG-BSA-Mg²⁺, NTG-BSA-Ca²⁺, NTG-BSA-Cu²⁺.




Fig 6: The Stern-Volmer plots of NTG quenching BSA fluorescence in the presence of different metal ions, (1) Cu²⁺, (2) Ca²⁺, (3) pure BSA, (4) Zn²⁺, (5) Mg²⁺.

Plots of $\log[(F_0-F)/F]$ versus $\log[Q]$ for NTG-BSA with different metal ions were showed in Fig 7. The value of K_a and binding sites (n) can be calculated by Eq. (1). The values of K_a for NTG associating with BSA in the presence of metal ions were presented in Table-4. The results showed that the values of binding constants of NTG to BSA were increased when Cu^{2+} and Mg^{2+} were titrated, but Ca^{2+} and Zn^{2+} decreased the values of binding constants. The number of binding site was about 1.0, which indicated that one binding site formed between NTG and BSA. The presence of Ca^{2+} , Zn^{2+} and Mg^{2+} , decreased the affinities of NTG to BSA and Cu^{2+} increased the affinities of NTG to BSA.




Fig 7: Double-logarithm curves of NTG quenching BSA fluorescence in the presence of metal ions, (1) Cu^{2+} , (2) Ca^{2+} , (3) pure BSA, (4) Zn^{2+} , (5) Mg^{2+}

Conclusions

Fluorescence and Ultraviolet-Visible methods for the determination of the interaction between NTG and BSA in the absence and presence of Cu^{2+} , Ca^{2+} , Mg^{2+} and Zn^{2+} was provided. NTG could quenching the fluorescence of BSA, the values of quenching rate constants suggested that the fluorescence quenching was a static process. The binding constant, the number of binding site, the thermodynamic parameters was significant to understand the mechanism of the drug with HSA in human. For many metal ions exist in organism, the study of drug-protein interaction in the presence of metal ions could provide many important information about the transport, absorption and distribution of drugs in plasma.

Acknowledgements

This work was supported by China Spark Program (grant numbers 2015GA690258); Jiangsu Province University Natural Science Foundation (grant numbers 14KJB180024); the open project of Jiangsu Key Laboratory for Bioresources of Saline Solis (grant numbers JKLBS2014009).

References

1. R. E. Olson and D. D. Christ, Plasma protein binding of drugs, *Annu. Rep. Med. Chem.*, **31**, 327 (1996).
2. B. Kaboudin, K. Moradi, M. R. Faghihi and F. Mohammadi, The fluorescence spectroscopic studies on the interaction of novel aminophosphinic acids with bovine serum albumin, *J. Lumin.*, **139**, 104 (2013).
3. S. Ghosh, S. Chakrabarty, D. Bhowmik, G. S. Kumar and N. Chattopadhyay, Stepwise unfolding of bovine and human serum albumin by an anionic surfactant: an investigation using the proton transfer probe norharmane, *J. Phys. Chem. B.*, **119**, 2090 (2015).
4. J. Maiti, S. Biswas, A. Chaudhuri, S. Chakraborty, S. Chakraborty and R. Das, Environment sensitive fluorescent analogue of biologically active oxazoles differentially recognizes human serum albumin and bovine serum albumin: Photophysical and molecular modeling studies, *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.*, **175**, 191 (2017).
5. N. Okabe and M. Hokaze, Effect of divalent metal ions on the binding of thyroxine to bovine serum albumin as measured by fluorescence, *Biol. Pharm. Bull.*, **16**, 719 (1993).
6. L. E. Wilkinson-White and S. B. Easterbrook-Smith, A dye-binding assay for measurement of the binding of Cu(II) to proteins, *J. Inorg. Biochem.*, **102**, 1831 (2008).
7. G. Navarra, A. Tinti, M. Leone, V. Militello and A. Torreggiani, Influence of metal ions on thermal aggregation of bovine serum albumin: aggregation kinetics and structural changes, *J. Inorg. Biochem.*, **103**, 1729 (2009).
8. K. C. Skyrianou, V. Psycharis, C. P. Raptopoulou, D. P. Kessissoglou and G. Psomas, Nickel-quinolones interaction. Part 4. Structure and biological evaluation of nickel(II)-enrofloxacin complexes compared to zinc(II) analogues, *J. Inorg. Biochem.*, **105**, 63 (2011).
9. P. N. Naik, S. A. Chimatadar and S. T. Nandibewoor, Interaction between a potent

- corticosteroid drug-Dexamethasone with bovine serum albumin and human serum albumin: A fluorescence quenching and fourier transformation infrared spectroscopy study, *J. Photochem. Photobiol. B.*, **100**, 147 (2010).
10. J. B. Xiao, T. T. Chen, L. S. Chen, H. Cao, F. Yang and Y. L. Bai, CdTe quantum dots (QDs) improve the affinities of baicalein and genistein for human serum albumin in vitro, *J. Inorg. Biochem.*, **104**, 1148 (2010).
 11. S. H. Cao, X. Y. Jiang and J. W. Chen, Effect of Zinc (II) on the interactions of bovine serum albumin with flavonols bearing different number of hydroxyl substituent on B-ring, *J. Inorg. Biochem.*, **104**, 146 (2010).
 12. T. Yamamoto, K. Shimizu, M. Takahashi, I. Tatsuno and K. Shirai, The effect of nitroglycerin on arterial stiffness of the aorta and the femoral-tibial arteries, *J. Atheroscler. Thromb.*, **24**, 1048 (2017).
 13. P. Derakhshan, S. H. Karbasy and R. Bahador, The effects of nitroglycerine on pain control during the propofol injection; a controlled, double-blinded, randomized clinical trial, *Anesth. Pain. Med.*, **5**, e26141 (2015).
 14. M. Goldin and G. A. Malanga, Tendinopathy: a review of the pathophysiology and evidence for treatment, *Phys. Sportsmed.*, **41**, 36 (2013).
 15. B. M. Andres and G. A. Murrell, Treatment of tendinopathy: what works, what does not, and what is on the horizon, *Clin. Orthop. Relat. Res.*, **466**, 1539 (2008).
 16. Y. Assem and M. Arora, Glyceryl trinitrate patches-An alternative treatment for shoulder impingement syndrome, *J. Orthopaedic. Translation.*, **3**, 12 (2015).
 17. A. R. Bokhari and G. A. C. Murrell, The role of nitric oxide in tendon healing, *J. Shoulder. Elbow. Surg.*, **21**, 238 (2012).
 18. E. D. Gambito, C. B. Gonzalez-Suarez, T. I. Oquiñena and R. B. Agbayani, Evidence on the effectiveness of topical nitroglycerin in the treatment of tendinopathies: a systematic review and meta-analysis, *Arch. Phys. Med. Rehabil.*, **91**, 1291 (2010).
 19. J. Wang, L. Liu, B. Liu, Y. Guo, Y. Zhang, R. Xu, S. Wang and X. Zhang, Spectroscopic study on interaction of bovine serum albumin with sodium magnesium chlorophyllin and its sonodynamic damage under ultrasonic irradiation, *Spectrochim. Acta A.*, **75**, 366 (2010).
 20. B. Tu, R. R. Li, Z. J. Liu, Z. F. Chen, Y. O. Yang and Y. J. Hu, Structure-activity relationship study between baicalein and wogonin by spectrometry, molecular docking and microcalorimetry, *Food Chem.*, **208**, 192 (2016).
 21. L. He, X. Wang, B. Liu, J. Wang, Y. Sun, E. Gao and S. Xu, Study on the interaction between promethazine hydrochloride and bovine serum albumin by fluorescence spectroscopy, *J. Lumin.*, **131**, 285 (2011).
 22. M. R. Eftink and C. A. Ghiron, Exposure of tryptophanyl residues in proteins. Quantitative determination by fluorescence quenching studies, *Biochemistry.*, **15**, 672 (1976).
 23. J. R. Lakowicz and G. Weber, Quenching of fluorescence by oxygen. A probe for structural fluctuations in macromolecules, *Biochemistry.*, **12**, 4161 (1973).
 24. L. Abdel-Aziz, L. Abdel-Fattah, A. El-Kosasy and M. Gaieda, A fluorescence quenching study of the interaction of nebivolol hydrochloride with bovine and human serum albumin, *J. appl. spectrosc.*, **82**, 620 (2015).
 25. S. Muzammil, Y. Kumar and S. Tayyab, Anion-induced stabilization of human serum albumin prevents the formation of intermediate during urea denaturation, *Proteins.*, **40**, 29 (2000).
 26. M. Pathak, R. Mishra, P.K. Agarwala, H. Ojha, B. Singh, A. Singh and S. Kukretic, Binding of ethyl pyruvate to bovine serum albumin: Calorimetric, spectroscopic and molecular docking studies, *Thermochim. Acta.*, **633**, 140 (2016).
 27. A. Ramanavicius, N. Ryskevici, A. Kausaite-Minkstimiene, U. Bubniene, I. Baleviciute, Y. Oztekin and A. Ramanaviciene, Fluorescence Study of Glucose Oxidase Self-Encapsulated Within Polypyrrole, *Sensor. Actuat. B-Chem.*, 171-172, 753-759 (2012).
 28. U. Bubniene, R. Mazetyte, A. Ramanaviciene, V. Gulbinas, A. Ramanavicius and R. Karpicz, Fluorescence Quenching Based Evaluation of Glucose Oxidase Composite with Conducting Polymer-Polypyrrole. *J. Phys. Chem C.*, **122**, 9491-9498 (2018).
 29. A. Ramanavicius, N. Ryskevici, Y. Oztekin, A. Kausaite-Minkstimiene, S. Jursenas, J. Baniukevic, J. Kirlyte, U. Bubniene and A. Ramanaviciene, Immunosensor based on fluorescence quenching matrix of conducting polymer-polypyrrole, *Anal. Bioanal. Chem.*, **398**, 3105-3113 (2010).
 30. X. Zhang, L. Li, Z. Xu, Z. Liang, J. Su, J. Huang and B. Li, Investigation of the interaction of naringin palmitate with bovine serum albumin: spectroscopic analysis and molecular docking, *Plos. One.*, **8**, e59106 (2013).
 31. P. D. Ross and S. Subramanian, Thermodynamics

- of protein association reactions: forces contributing to stability, *Biochemistry*, **20**, 3096 (1981).
32. M. A. Rub, J. M. Khan, A. M. Asiri, R. H. Khan and K. Din, Study on the interaction between a amphiphilic drug and bovine serum albumin: A thermodynamic and spectroscopic description, *J. Lumin.*, **155**, 39 (2014).
33. Y. J. Hu, Y. Liu, L. X. Zhang, R. M. Zhao and S. S. Qu, Studies of interaction between colchicine and bovine serum albumin by fluorescence quenching method, *J. Mol. Struct.*, **750**, 174 (2005).
34. L. Z. Wu, B. L. Ma, D. W. Zou, Z. X. Tie, J. Wang and W. Wang, Influence of metal ions on folding pathway and conformational stability of bovine serum albumin, *J. Mol. Struct.*, **877**, 44 (2008).
35. M. Karahan, Z. Mustafaeva and C. Ozeroğlu, Investigation of ternary complex formations of polyacrylic acid with bovine serum albumin in the presence of metal ions by fluorescence and dynamic light scattering measurement, *Protein J.*, **29**, 336 (2010).
36. Z. Jing, R. Qi, C. Liu and .P. Ren, Study of interactions between metal ions and protein model compounds by energy decomposition analyses and the AMOEBA force field. *J. Chem. Phys.*, **147**, 161733 (2017).
37. Y. Zhang, S. Shi, K. Huang, X. Chen and M. Peng, Effect of Cu^{2+} and Fe^{2+} for drug delivery: Decreased binding affinity of ilaprazole to bovine serum albumin, *J. Lumin.*, **131**, 1927 (2011).