

## Synthesis, Crystal Structure and Antifungal Activity of 1-(3,3-Dimethyl-2-oxobutyl)-N-phenyl-1H-1,2,4-triazole-3-carboxamide

<sup>1</sup>Guang Rong Cai<sup>§\*</sup>, <sup>2</sup>Zhi Fang Li<sup>§</sup>, <sup>1</sup>Yan Chuan Gong, <sup>1</sup>Xue Dong and <sup>1</sup>Wen Su Cai  
<sup>1</sup>Sichuan Provincial Key Laboratory for Characteristic Plant Development and Research,  
Sichuan University of Arts and Science, Dazhou, 163000, China.  
<sup>2</sup>Dazhou Vocational College of Chinese Medicine, Dazhou, 163000, China.  
[swueducaai@sina.com](mailto:swueducaai@sina.com)\*

§These authors contributed equally this work.

(Received on 24<sup>th</sup> July 2023, accepted in revised form 17<sup>th</sup> January 2024)

**Summary:** Fungicides play a vital role in protecting crops from fungal damage. However, fungicide resistance is one of the most important issues in modern agriculture. Hence, it is very necessary to develop new fungicides continuously. Triazole compounds have received considerable interest in agricultural chemistry due to a novel action mode, extremely high activity against phytopathogenic fungi, low acute toxicity to mammals, and environmentally benign characteristics. The title compound 1-(3,3-dimethyl-2-oxobutyl)-N-phenyl-1H-1,2,4-triazole-3-carboxamide 5, synthesized using methyl 1H-1,2,4-triazole-3-carboxylate 1 as the start material, was successfully obtained via multiple synthesis route and finally characterized by <sup>1</sup>H NMR, <sup>13</sup>C NMR, HRMS and single-crystal X-ray diffraction. Compound 5 (C<sub>15</sub>H<sub>18</sub>N<sub>4</sub>O<sub>2</sub>, Mr = 287.1500) belongs to the orthorhombic system, space group Pn2<sub>1</sub>a, with a = 14.49451(11) Å, b = 20.34686(18) Å, c = 10.17021(9) Å, V = 2999.38(4) Å<sup>3</sup>, Z = 8, D<sub>c</sub> = 1.268 g/cm<sup>3</sup>, T = 293.55(14) K, μ (CuKα) = 0.710 mm<sup>-1</sup>, F(000) = 1216.0, the final R = 0.0448, and wR = 0.1260 with I > 2σ (I). Furthermore, the results from biological assays indicated that the title compound exhibited a similar antifungal activity (EC<sub>50</sub> = 8.98 mg·L<sup>-1</sup>) compared to triadimefon (EC<sub>50</sub> = 5.6 mg·L<sup>-1</sup>) against *G. cingulata*. And had different degrees of weak activity against other phytopathogenic fungi, including *A. solani*, *S. sclerotiorum*, *G. saubinetii* and *T. cucumeris*. Potentially, the result lay the foundation for the development of novel fungicides.

**Keywords:** Synthesis; Crystal structure; Triazole; Amide; Antifungal activity

### Introduction

Fungicides, a major part of crop protection products, widely applied to major crops such as rice, corn and wheat, are playing an essential role by increasing both crop quality and yield [1]. Triazole fungicides belong to the sterol demethylation inhibitor (DMI) group, which is characterized by the inhibition of ergosterol biosynthesis, an important component of the fungal cell membrane [2], have been extensively and widely used in wheat, rice and corn fields for managing fungal diseases, including leaf rust, powdery mildew, leaf spots gibberellic diseases and sheath blight [3-4]. However, in recent years the developments of resistance to currently available triazole-antifungicide have been reported around the world [5-7]. Fungal diseases management is still a challenge for researchers, and continuous innovation is essential to maintain the effectiveness of pathogen management.

1,2,4-triazole derivatives are often used in the medicinal and agricultural areas. Some of them had been developed as commercial pesticides, such as Diniconazole [8], Triadimefon [9], Triadimenol [10], Flusilazole [11], Epoxiconazole [12], Prothioconazole

[13] and Tebuconazole [14]. Also, tert-butyl ketone structural element displayed outstanding activity in agrochemical, such as fungicidal activity [15-16], phytohormone activity [17-18]. Meanwhile, Amide fungicides are widely used for the highly effective fungicidal activity, and performed outstanding effects on downy mildew on soybean, cucumber, millet, potato and eggplant caused by *Phytophthora* [19]. Unfortunately, after using for a period of time, the fungicides will be gradually out of service [20-22]. Therefore, it is urgent to develop some innovative compounds displaying excellent fungicidal activity.

In view of these facts mentioned above, the title compounds were designed by introducing tert-butyl ketone and amide pharmacophore into 1,2,4-triazole scaffold. A new 1,2,4-triazole derivative was synthesized and characterized by <sup>1</sup>H NMR, <sup>13</sup>C NMR and HRMS. The single-crystal structure of the title compound was determined by X-ray diffraction. The antifungal activity of the title compound was investigated.

---

\*To whom all correspondence should be addressed.


## EXPERIMENTAL

## General Techniques

1*H*-1,2,4-triazole-3-carboxylate, 1-chloro-3,3-dimethyl-butan-2-one, aniline and 2-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU), Potassium carbonate ( $K_2CO_3$ ) and polyethylene glycol-600 (PEG-600) were purchased from Shanghai Aladdin Biochemical Technology Co., Ltd. (Shanghai, China). Unless otherwise noted, all reagents were obtained from commercial suppliers and were used without further purification. All reactions were monitored by TLC with GF254 silica gel plates. Flash column chromatography was carried out using 200-300 mesh silica gel by using certain ratio of eluent with petroleum ether/ethyl acetate/methanol. Single crystal of compound **5** was cultured in a system of dichloromethane and petroleum ether. The melting point was measured on a Hanon MP100 automatic melting point apparatus (Jinan Hanon Instruments Co., Ltd., Jinan, Shandong, China), using an open capillary tube. The NMR spectra was recorded at 400 MHz ( $^1H$  NMR) and 100 MHz ( $^{13}C$  NMR) on a Bruker AV-400 spectrometer with TMS as an internal standard. The chemical shifts ( $\delta$ ) are given in ppm, and the coupling constants ( $J$ ) in Hz. High-resolution mass spectral analysis was carried out on an FTICR-MS Varian 7.0 T FTICR-MS instrument (Varian IonSpec, Lake Forest, CA, USA). Single-crystal X-ray structure was measured on a Bruker SMART APEX II X-ray single crystal diffractometer (Bruker AXS, Karlsruhe, BW, Germany).

## Synthetic Procedure

The synthetic route of Compound **5** is outlined in Scheme 1. The intermediates **3** and **4** were synthesized according to the reference [23]. A mixture of 1-(3,3-dimethyl-2-oxobutyl)-1*H*-1,2,4-triazole-3-carboxylic acid



Scheme-1: Synthesis route of compound **5**. Regent and condition: (i) AcOEt,  $K_2CO_3$ -PEG (600), 56 °C, 78 %; (ii) NaOH (10 %), MeOH:  $H_2O$  (V/V) = 1:1, 30 min, rt,  $\geq 99\%$ ; (iii) Aniline, DMC, HATU, TEA, r.t, 61 %.

**4** (506.4 mg, 2.4 mmol), triethylamine (600  $\mu$ L) and HATU (1 eq) in dichloromethane (50 mL) was stirred at r.t for 1 h. Afterwards, aniline (232.8 mg, 2.4 mmol) was added and stirring was continued for 40 min at same temperature. The reaction mixture was washed with water (310 mL) and saturated salt water (30 mL), dried over anhydrous  $MgSO_4$ , filtered through a Celite pad (i.d. = 8 cm; height = 1.5 cm) and concentrated under reduced pressure. The crude product was purified by flash chromatography (petroleum ether / ethyl acetate/ methanol 3:2:0.5) to provide compound **5** (420 mg, 61%) as white powder, m.p. 152-154°C;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.96 (s, 1H), 8.22 (s, 1H), 7.71 (d,  $J = 7.7$  Hz, 2H), 7.37 (t,  $J = 7.9$  Hz, 2H), 7.16 (d,  $J = 7.4$  Hz, 1H), 5.32 (s, 2H), 1.27 (s, 9H).  $^{13}C$  NMR (100 MHz,  $CDCl_3$ )  $\delta$  205.76, 156.89, 156.57, 145.63, 137.33, 129.12, 124.67, 119.90, 54.08, 43.58, 26.14. HR-ESI-MS for (ESI $^+$ ) for  $C_{15}H_{18}N_4O_2$  (m/z): calcd. 287.1463, found: 287.1500[M+H] $^+$ .

## Structure Determination

Single crystals of Compound **5** were obtained by recrystallization from a system of dichloromethane and petroleum ether at room temperature. The crystal dimensions were 0.20  $\times$  0.05  $\times$  0.05 mm $^3$ . The reflection data of Compound **5** was collected by using X-radiation ( $\lambda = 1.54184$  Å) at 293(14) K via a Bruker SMART APEX II X-ray single crystal diffractometer (Bruker AXS, Karlsruhe, BW, Germany). A total of 30970 reflections were collected, 5813 of which were independent with  $R_{int} = 0.0239$ . The structure of Compound **5** was solved via a direct method using SHELXS-97 (University of Gottingen, Gottingen, NI, Germany). The solutions were refined by full-matrix least squares techniques on  $F^2$  by SHELXL-2015 program [24]. The final cycle of refinement gave  $R = 0.0461$  and  $wR = 0.1282$ . Selected crystallographic data of the Compound **5** is provided in Table 1.

Table-1: Selected crystallographic data of Compound 5.

Parameter	Result	Parameter	Result
CCDC No.	2088260	$\alpha/^\circ$	90
Empirical formula	$C_{15}H_{18}N_4O_2$	$\beta/^\circ$	90
Formula weight	286.33	$\gamma/^\circ$	90
Temperature/K	293.55(14)	Volume/ $\text{\AA}^3$	2999.38(4)
Crystal system	orthorhombic	Z	8
Space group	$Pn2_1a$	$\rho_{\text{calc}}/\text{cm}^3$	1.268
a/ $\text{\AA}$	14.49451(11)	Reflections collected	30970
b/ $\text{\AA}$	20.34686(18)	Independent reflections	[ $R_{\text{int}} = 0.0239$ , $R_{\text{sigma}} = 0.0148$ ]
c/ $\text{\AA}$	10.17021(9)	Data/restraints/parameters	5813/1/385
$\mu/\text{mm}^{-1}$	0.087	Goodness-of-fit on $F^2$	1.039
F(000)	1216.0	Final R indexes [ $I \geq 2\sigma(I)$ ]	$R_1 = 0.0448$ , $wR_2 = 0.1260$
Crystal size/ $\text{mm}^3$	$0.2 \times 0.05 \times 0.05$	Final R indexes [all data]	$R_1 = 0.0461$ , $wR_2 = 0.1282$
Radiation	$\text{CuK}\alpha$ ( $\lambda = 1.54184$ )	Largest diff. peak/hole / $e \text{\AA}^{-3}$	0.38/-0.19
2 $\theta/^\circ$	8.692 to 143.53		
Index ranges	$-17 \leq h \leq 12$ , $-25 \leq k \leq 24$ , $-12 \leq l \leq 12$		

### Antifungal Activity

The antifungal activities of Compound 5, Triadimefon were tested in vitro against *Alternaria solani*, *Sclerotinia sclerotiorum*, *Gibberella saubinetii*, *Glomeran cingulata* and *Thanatephorus cucumeris*. The five kinds of test fungi were purchased from National Microbial Resource Center (Beijing, China). After retrieval from the storage tube, the strains were incubated on potato dextrose agar (PDA) at 27 °C for 4 days to get new mycelia for the antifungal tests. The fungicidal activity of the target compound was tested in vitro against the five plant pathogenic fungi using the mycelia growth inhibition method [25]. The tested compound were dissolved in DMSO to prepare a 1000  $\text{mg.L}^{-1}$  stock solution before mixing with PDA. The media containing compounds at a concentration of 50  $\text{mg.L}^{-1}$  were then poured into sterilized Petri dishes for initial screening. After two days at 27 °C, the colony diameter of each strain was measured. Percentage inhibition rate was calculated as  $(1-a/b) \times 100\%$ , where a represents the colony diameter in the Petridishes with tested compounds and b is the mean colony diameter in control Petri dishes. Each test was repeated three times. The 1000  $\text{mg.L}^{-1}$  solution was diluted to 200, 100, 50, 25, 12.5, 6.25, 3.12, 1.56, 0.78  $\text{mg.L}^{-1}$  and the above experiments were repeated, the inhibition rates were calculated separately. The  $\text{EC}_{50}$  values were calculated using SPSS Statistics v 22.0.

### Result and Discussion

#### Description of the Structure

Compound 5 crystallized in the orthorhombic system. The  $Pn2_1a$  space group and the molecular structure of Compound 5 are depicted in Figure 1. Selected molecular structure parameters (bond lengths and bond angles) for Compound 5 can be found summarized in Table 1. The packing arrangement is shown in Figure 2.

The crystal data for Compound 5 was deposited at the Cambridge Crystallographic Data Centre (12 Union Road, Cambridge CB2 1EZ, UK; fax: +44-1223-336033; e-mail: deposit@ccdc.cam.ac.uk) as supplementary publication No. CCDC-2088260. Crystallographic data for this crystal is available free of charge at the following website: [http://www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif) or from the Cambridge Crystallographic Data Centre.




Fig. 1: Crystal structure of Compound 5.

The bond lengths and bond angles of the aromatic rings (phenyl and triazolyl) in this crystal structure are in accordance with the general normal ranges [26-27]. The phenyl ring and triazolyl ring were both connected by an amide group, and the  $\text{C}(9)=\text{O}(2)$  bond length in this amide group was 1.219(3)  $\text{\AA}$ . Another carbonyl ( $\text{C}(1)=\text{O}(5)$ ) bond length was 1.202(4). The both is shorter to the general  $\text{C}=\text{O}$  double-bond length reported in the literature [ 28-29]. Compared to the normal  $\text{C}-\text{N}$  bond (1.47~1.50  $\text{\AA}$ ), the shorter  $\text{C}(7)-\text{N}(1)$  (1.331(2)  $\text{\AA}$ ),  $\text{C}(7)-\text{N}(3)$  (1.322(4)  $\text{\AA}$ ) and  $\text{C}(8)-\text{N}(2)$  (1.320(3)  $\text{\AA}$ ) bonds in the triazole ring,  $\text{C}(9)-\text{N}(4)$  (1.351(4)  $\text{\AA}$ ) of amide and  $\text{C}(10)-\text{N}(4)$  (1.421(4)  $\text{\AA}$ ) may be due to the effect of conjugation. The bond angles of triazole ring ( $\text{N}(1)/\text{N}(2)/\text{C}(7)/\text{N}(3)/\text{C}(8)$ ) varied from  $102.0(2)^\circ$  to  $115.3(2)^\circ$  with the average of  $108^\circ$ , and benzene ring ( $\text{C}(10)/\text{C}(11)/\text{C}(12)/\text{C}(13)/\text{C}(14)/\text{C}(15)$ ) varied from  $119.6(3)^\circ$  to  $120.2(3)^\circ$  with the average of  $120^\circ$ .




Fig. 2: Packing arrangement of Compound 5 in the unit cell.




Fig. 3: Crystal unit stacking of compound 5.

Table-2. EC<sub>50</sub> values of compound 5, triadimefon against *Alternaria solani*, *Sclerotinia sclerotiorum*, *Gibberella saubinetii*, *Gloneman cingulatal* and *Thanatephorus cucumeris*<sup>1</sup>.

	EC <sub>50</sub> (± SD) mg.L <sup>-1</sup>				
	<i>A. solani</i>	<i>S. sclerotiorum</i>	<i>G. saubinetii</i>	<i>G. cingulatal</i>	<i>T. cucumeris</i>
Compound 5	125.24±1.786	78.22±1.334	59.98±2.311	8.98±0.484	66.20±1.681
Triadimefon	1.05±0.922	8.45±0.264	19.35 ±0.459	5.6±0.504	16.63±0.162

The crystal packing characteristics of Compound **5** in the unit cell are described in Figure 2. Two adjacent molecules with a head to tail arrangement (Figure 3) are found in the crystal packing. In the intermolecular, it is worth noting that face-to-face  $\pi$ - $\pi$  stacking pattern of the title compound, which would be proved by the relative positions between triazole ring and phenyl ring of the two molecules: the centroid separation of them is 3.5696 Å, and their angle is 6.9547°. These interactions are estimated to play a role in stabilizing the crystal structure.

#### Spectroscopic Properties

The structure of compound **5** was confirmed via melting point,  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, and HRMS analysis. Signals corresponding to the C-H proton in the triazole ring and N-H proton in the amide group were observed at  $\delta$  8.22 and  $\delta$  8.96, respectively. The signals corresponding to the protons on the benzene ring were observed at  $\delta$  7.16-7.71. The HRMS data of compound **5** was in good agreement with the theoretical data that was calculated on the basis of the molecular formula.

#### Evaluation of Antifungal Activity

The *in vivo* fungicidal activities of the compound **5** against *Alternaria solani*, *Sclerotinia sclerotiorum*, *Gibberella saubinetii*, *Gloneman cingulatal*, and *Thanatephorus cucumeris* were evaluated, and triadimefon was used as controls (Table-2). The primary bioassay showed the title compound exhibits weak inhibiting activity towards *Alternaria solani*, *Sclerotinia sclerotiorum*, *Gibberella saubinetii*, and *Thanatephorus cucumeris* with  $\text{EC}_{50}$  of 125.24 mg. L<sup>-1</sup>, 78.22 mg. L<sup>-1</sup>, 59.98 mg. L<sup>-1</sup>, and 66.20 mg.L<sup>-1</sup>, respectively. Interestingly, the inhibitory activity of title compound against *Gloneman cingulatal* was close that of triadimefon. These results indicated that compound **5** could be further used as a lead compound to develop novel fungicides.

<sup>1</sup>A. *solani*: *Alternaria solani*; *S. sclerotiorum*: *Sclerotinia sclerotiorum*; *G. saubinetii*: *Gibberella saubinetii*; *G. cingulatal*: *Gloneman cingulatal* and *T. cucumeris*: *Thanatephorus cucumeris*. The experiment was carried out in three triplicates. The data of the fungicidal activities were statistically analyzed using the SPSS 22.0 software package to obtain  $\text{EC}_{50}$  values. The latter represent the mean  $\pm$  standard deviation (SD) of triplicate experiments.

#### Conclusions

In summary, the compound 1-(3,3-dimethyl-2-oxobutyl)-N-phenyl-1H-1,2,4-triazole-3-carboxamide **5** was synthesized and characterized by  $^1\text{H}$

NMR,  $^{13}\text{C}$  NMR, HRMS, and X-ray diffraction. The synthesis followed a strategy of inserting an amide group between a benzene ring and a triazole ring. The biological assay results indicated compound **5** showed a similar antifungal activity compared to triadimefon against *Gloneman cingulatal*. Potentially, the results obtained will lay the foundation for the design and development of novel fungicides.

#### Acknowledgments

We are grateful for financial support by the Field Grade Foundation of Si Chuan University of Arts and Science (No.2019PT004Z) ; the Key Laboratory of Sichuan University for the Development and Research of Characteristic Plants Open Fund Project (No.TSZW2001) and Sichuan University of Arts and Sciences Doctoral Research Start-up Fund (2021RC002Z) .

#### References

1. M. Y. Yoon, B. Cha, J. C. Kim, Recent trends in studies on botanical fungicides in agriculture, *Plant. Pathol. J.*, **29**, 1 (2013).
2. Y. Tatsumi, M. Nagashima, T. Shibanushi, A. Iwata, Y. Kangawa, F. Inui, Y. Nishiyama, Mechanism of action of efinaconazole, a novel triazole antifungal agent, *Antimicrob Agents. Chemother.*, **57**, 2405 (2013).
3. R. H. Bromilow, A. A. Evans, P. H. Nicholls, Factors affecting degradation rates of five triazole fungicides in two soil types: 2. Field studies, *Pestic. Sci.*, **55**, 1135 (1999).
4. M. Beyer, M. B. Klix, H. Klink, J. A. Verreet, Quantifying the effects of previous crop, tillage, cultivar and triazole fungicides on the deoxynivalenol content of wheat grain—a review, *J. Plant. Dis. Protect.*, **113**, 241 (2006).
5. P. E. Verweij, G. H. Kema, B. Zwaan, W. J. Melchers, Triazole fungicides and the selection of resistance to medical triazoles in the opportunistic mould *Aspergillus fumigatus*, *Pest. Manag. Sci.*, **69**, 165 (2013).
6. E. Snelders, S. M. Camps, A. Karawajczyk, G. Schaftenaar, G. H. Kema, H. A. Van der Lee, P. E. Verweij, Triazole fungicides can induce cross-resistance to medical triazoles in *Aspergillus fumigatus*, *PloS. One.*, **7**, 31801 (2012).
7. O. Bader, J. Tünnermann, A. Dudakova, M. Tangwattanachuleeporn, M. Weig, U. Groß, Environmental isolates of azole-resistant *Aspergillus fumigatus* in Germany. *Antimicrob. Agents, Chemother.*, **59**, 4356 (2015).

8. M. J. Jung, H. Y. Kim, K. B. Lim, Effect of diniconazole on the growth and taking roots after transplanting of *Sesamum indicum* 'Baekseol' plug seedlings, *J. Crop. Sci. Biotech.*, **23**, 235 (2020).
9. C. A. Jaleel, R. Gopi, R. Panneerselvam, Growth and photosynthetic pigments responses of two varieties of *Catharanthus roseus* to triadimefon treatment, *Comptes. Rendus Biologies.*, **331**, 272 (2008).
10. J. P. Da Silva, A. M. Da Silva, I. V. Khmelinskii, J. M. G. Martinho, L. V. Ferreira, Photophysics and photochemistry of azole fungicides: triadimefon and triadimenol, *J. Photoch. Photobio. A.*, **142**, 31 (2001).
11. D. U. Ozakca, H. Silah, Genotoxicity effects of Flusilazole on the somatic cells of *Allium cepa*, *Pestic. Biochem. Physiol.*, **107**, 38 (2013).
12. Y. X. Wu, A. Von Tiedemann, Physiological effects of azoxystrobin and epoxiconazole on senescence and the oxidative status of wheat, *Pestic. Biochem. Physiol.*, **71**, 1 (2001).
13. K. Audenaert, E. Callewaert, M. Höfte, S. De Saeger, G. Haesaert, Hydrogen peroxide induced by the fungicide prothioconazole triggers deoxynivalenol (DON) production by *Fusarium graminearum*, *Bmc. Microbiol.*, **10**, 1 (2010).
14. P. A. Paul, P. E. Lipps, D. E. Hershman, M. P. McMullen, M. A. Draper, L. V. Madden, A quantitative review of tebuconazole effect on *Fusarium* head blight and deoxynivalenol content in wheat, *Phytopathology.*, **97**, 211 (2007).
15. A. K. Jha, P. K. Shukla, N. Soni, A. Verma, Synthesis, characterization and biological evaluation of thiazole incorporated triazole compounds, *Der Pharmacia Lettre.*, **7**, 67 (2015).
16. W. Wang, C. Sheng, X. Che, H. Ji, Z. Miao, J. Yao, Design, synthesis, and antifungal activity of novel conformationally restricted triazole derivatives. *Archiv. Der. Pharmazie.*, **342**, 732 (2010).
17. M. Zhang, L. Duan, X. Tian, Z. He, J. Li, B. Wang, Z. Li, Uniconazole-induced tolerance of soybean to water deficit stress in relation to changes in photosynthesis, hormones and antioxidant system, *J. Plant. Physiol.*, **164**, 709 (2007).
18. M. Berova, Z. Zlatev, Physiological response and yield of paclobutrazol treated tomato plants (*Lycopersicon esculentum* Mill.), *Plant. Growth. Regul.*, **30**, 117 (2000).
19. P. Huang, S. Zhou, Y. Du, H. Li, Y. Lv, L. Lv, Study on a new type of high efficient amide compound fungicides against soybean rust, *Tetrahedron. Lett.*, **64**, 152745(2021).
20. M. Blum, M. Waldner, G. Olaya, Y. Cohen, U. Gisi, H. Sierotzki, Resistance mechanism to carboxylic acid amide fungicides in the cucurbit downy mildew pathogen *Pseudoperonospora cubensis*, *Pest. Manag. Sci.*, **67**, 1211 (2011).
21. D. Fernández-Ortuño, A. Grabke, X. Li, G. Schnabel, Independent emergence of resistance to seven chemical classes of fungicides in *Botrytis cinerea*, *Phytopathology.*, **105**, 424 (2015).
22. P. Bowyer, D. W. Denning, Environmental fungicides and triazole resistance in *Aspergillus*, *Pest. Manag. Sci.*, **70**, 173 (2014).
23. G. R. Cai, G. G. Zuo, D. F. Zheng, N. J. Feng, Synthesis and Biological Evaluation of 3, 3-Dimethyl-1-(1 H-1, 2, 4-triazole-1-yl) butan- 2-One Derivatives as Plant Growth Regulators, *Chem. Res. Chinese U.*, **35**, 221 (2019).
24. G. M. Sheldrick, SHELXT-Integrated space-group and crystal-structure determination, *Acta. Crystallogr. A.*, **71**, 3 (2015).
25. J. X. Mu, Z. W. Zhai, M. Y. Yang, Z. H. Sun, H. K. Wu, X. H. Liu, Synthesis, crystal structure, DFT Study and antifungal activity of 4-(5-((4-bromobenzyl) thio)-4-phenyl-4H-1, 2, 4-triazol-3-yl) pyridine, *Crystals.*, **6**, 4 (2016).
26. K. Peters, E. M. Peters, T. Irrgang, A. Hetzheim, Crystal structure of 3,4-diphenyl-6- methyl-1-propargylaminoimidazo[1,5-b]pyridazine, *Zeitschrift fur Kristallographie-New Crystal Structures.*, **214**, 345 (1999).
27. R. Brehme, E. Gründemann, M. Schneider, R. Radeaglia, G. Reck, B. Schulz, Aza-enamines X: Formylation of pyrazole-4-carbaldehyde hydrazones at the hydrazonoazomethine C-atom. *Cheminform*, **10**, 1615(2003).
28. J. Q. Weng, L. Wang, X. H. Liu, Synthesis, crystal structure and herbicidal activity of a 1,2,4-triazol-5-(4H)-one derivative, *J. Chem. Soc. Pak.*, **34**, 1248 (2012).
29. R. Löser, R. Pitzschler, M. Köckerling, R. Löser, R. Pitzschler, M. Köckerling, Synthesis and X-ray crystal structure of N'-cyano-N,N' -dimethyl-4-nitrobenzohydrazide, *Crystals*, **7**, 290 (2017).