

Documentation of Ethnomedicinal Uses of Wild Plants Growing in Kodo Mountain by Kurdish Tribe of Iraq

¹Faiq H.S. Hussain*, ¹Soma Majedi, ^{1,2}Tola Abdulsattar Faraj, ³Mehmet Ozdemir, ⁴Javed Ahamad, ⁵Tariq H. Karim, ¹Kovan Dilawer Issa and ¹Mohammad Q. Mustafa
¹Department of Medical Analysis, Faculty of Applied Science, Tishk International University – Erbil 44001
 Kurdistan Region, Iraq.
²Department of Basic Sciences, College of Medicine, Hawler Medical University
 Erbil 44001, Kurdistan Region, Iraq.
³Department of Dentistry, Faculty of Dentistry, Tishk International University
 Erbil 44001, Kurdistan Region, Iraq.
⁴Department of Pharmacognosy, Faculty of Pharmacy, Tishk International University
 Erbil 44001, Kurdistan Region, Iraq.
⁵Civil Engineering Department, Faculty of Engineering, Tishk International University
 Erbil 44001, Kurdistan Region, Iraq.
 faiq.hussain@tiu.edu.iq*

(Received on 4th May 2023, accepted in revised form 8th November 2023)

Summary: Medicinal plants have a long history of different uses and are still of great importance in the daily life of the Kurds living in the Kurdistan province of Iraq. The present review provides comprehensive phytochemical and pharmacological information about medicinal plants growing in the Kodo Mountain area. In fact, no systematic study of the bioresources from this unique region has yet been reported in the scientific literature, even if local villagers have been consuming several plants as nutrition and ethnomedicinal food for centuries, until today.

The data reported in this paper were obtained through interviews with local herbal healers and people living in villages around the Kodo Mountains. They concern 40 plants belonging to 16 botanical families, which are considered medicinal by the local population. The present study recorded and examined the demographic information given by the study participants, the local names and the parts used of the plants, the preparation and administration techniques, and the treatments for diseases. The most frequently cited families were Asteraceae (32.5 %), Lamiaceae (10 %), Brassicaceae, Malvaceae, and Papaveraceae (7.5 % each). The main parts of the plants used for medicinal purposes were leaves (36 %) and flowers (29.5 %). They were administered as decoctions (42 %), raw (26 %), or powder (14 %). Several important phytochemicals have been isolated from the used plants, including flavonoids (60 %), terpenoids (45 %), phenolic acids (42.5 %), polyphenolic compounds (40 %), and essential oils (30 %). The plant ethnomedicinal and pharmacological uses were supported by their antibacterial (18%, Apiaceae, Lamiaceae, Papaveraceae), anti-inflammatory (18%, Malvaceae, Asteraceae, Papaveraceae), anti-oxidant (16%, Malvaceae, Apiaceae, Lamiaceae, Asteraceae, Papaveraceae), anti-cancer (9%, Lamiaceae, Papaveraceae, Asteraceae), anti-parasitic (8%, Asteraceae), hepatoprotective (7%, Asteraceae, Lamiaceae, Papaveraceae, Fabaceae), anti-diabetic (5%, Asteraceae, Fabaceae), anti-fungal (5%, Amaryllidaceae, Umbelliferae, Asphodelaceae, Orchidaceae), anti-spasmodic (4%, Asteraceae, Papaveraceae), and diuretic (3%, Asteraceae, Amaryllidaceae, Plantaginaceae) activities. This study illustrates the significance of traditional medicinal plants that have been utilized for treatment and healing the wounds and curing the illnesses Kurdish tribe in Kodo mountain, north-east part of Kurdistan Region in Iraq, that can be used as reference for further investigations for the researchers in future.

Keywords: Kodo Mountain, Phytochemistry, Ethnopharmacology, Medicinal plants, Metabolites.

Introduction

Ethnobotany can be defined as the study of the relationship between people and plants. The term "ethnobotany" was coined in 1896 by the American botanist John Harshberger (1) to describe the study of plants used by the peoples living in a particular area, considering the cultural and the ecological contexts in which they are. For the entire history of humanity, especially medicinal plants have been used

extensively worldwide to treat a wide range of disorders and maladies and have had a significant role in international trade. Peoples have learnt how to use medicinal herbs by trial and error, and knowledge has usually been passed from one generation to another only orally. The geographical location of the Iraqi Kurdistan Region, and the geological and ecological

*To whom all correspondence should be addressed.

features make the region's biodiversity one of the richest in Iraq (2).

The Kodo Mountain is located about 103 km north-east of Erbil, capital of the Kurdistan Region (Fig 1). The flora and the ethnobotany of the Kodo Mountain region have not yet been fully investigated. Therefore, we started this work with the aim of raising awareness of the region's botanical resources that need to be valued and protected. In particular, we focused

on the possible relationship between traditional applications and medicinal effects, and we evaluated the therapeutic potential of the region's natural remedies. Moreover, the information given in this study adds to the limited number of scientific studies recently conducted about the structures and bioactivities of metabolites isolated from medicinal plants native to regions near the Kodo Mountain (2-12).




Fig 1: A. Kodo Mountain, Kurdistan region, Erbil–Iraq. B. Map of Kurdistan Region–Iraq.



Fig. 2: Interviews with local people living around the Kodo Mountain areas.

Plant Material

A total of 40 medicinal plants (Table 1) were mentioned by local people in the interviews. Samples of the plants (Fig 3) were collected and identified by Prof. Abdul-Hassan Al-Khayat at Salahaddin University and are deposited at Applied Science

Faculty Herbarium, Tishk International University, Erbil, Kurdistan Region of Iraq.

The families of Asteraceae, Lamiaceae, Apiaceae, Malvaceae, and Papaveraceae included the greatest number of plants, making the 65% of the cited species (Fig 4).




Fig 3: Some Plants Growing in the Kodo Mountain Area, A: *Dactylorhiza umbrosa*, B: *Achillea ageratum*, C: *Artemisia absinthium*, D: *Astragalus glaucacathus*, E: *Asphodelus albus*, F: *Cichorium intybus*.




Fig 4: Percent distribution of the botanical families of plants collected on the Kodo Mountain.

Phytochemical Aspects

A search in the scientific literature revealed that more than 150 compounds were isolated and identified from the plants listed in Table 1. Flavonoids (60%) were the compounds most isolated from the plants, followed by terpenoids (45%) and phenolic acids (42.5%). Other compounds which were also present in significant amounts are polyphenolic derivatives (40%), essential oils (30%), alkaloids (27.5%), carbohydrates (27.5%), steroids (25%) and fatty acids (20%), while the remaining compounds occurred in minor amounts (Fig 5).

The plants were most consumed as decoction (42%) or as raw (26%). Less frequently, they were consumed as a powder (14%), an infusion (8%), a chew?? (4%), a gum, a juice, and by smoking (2% each)

The most used parts of plants were leaves (36%) and flowers (29.5%), followed by roots (8%), seeds and stems (6% each), fruits and bulbs (each 5% each), aerial parts (3%), and latex (1.5%).

Biological and Pharmacological Aspects

Over the whole history of human life plants have supplied food, various types of flavors,

medications, garments, and protection means worldwide. Nowadays, the study of outstanding medicinal properties and application of plants native to different regions is one of the most active research fields. The great biodiversity of plants growing on the slopes of the Kodo Mountain and the lack of a systematic scientific knowledge prompted us to conduct a comprehensive study (Table 1). Indeed, the interviews with the people living in this rural area demonstrated that almost all native plants have some applications; above all, they are used as traditional remedies to treat most diseases, except acute and urgent cases that require a surgery. Anti-inflammatory, antibacterial, anticancer, cardioprotective, antioxidant, and antiparasitic effects are the properties most frequently attributed to the plants growing in the Kodo Mountain (Fig 6). Each group of plants displays a characteristic range of biological activities, as shown in Fig 7 for the families containing the highest numbers of species. The families of Asteraceae and Papavareaceae exhibited the widest spectra of pharmacological effects, while Malvaceae showed only two types of biological properties. Within the Asteraceae, *Achillea ageratum*, *A. millefolium*, and *A. santolina* are examples of plants widely used by local herbalist, which exhibited many beneficial effects (Table 1).




Fig. 5: Percentages of the most abundant natural products isolated from the investigated plants and families.




Fig. 6: Percentages of biological activities attributed to plants collected in the Kodo Mountain.

Thus, *Achillea ageratum* has showed antioxidant, cytotoxic, antibacterial, and anti-inflammatory properties. On the other hand, antispasmodic, anticancer, hepatoprotective, antidiabetic, and antileishmanial activities have been determined for the flower and leaf extracts of *A.*

millefolium and *A. santolina*. *Anthemis des champs* (*A. arvensis*), *Artemisia absinthium*, *Cota tinctoria*, *Echinops viscosus nozaff* are other species in the Asteraceae family that possess remarkable pharmacological effects (Table 1).




Fig 7: Percentages of various biological activities determined for the five families containing the highest number of species.

Table-1: Literature data concerning medicinal plants growing in Kodo Mountain.

S.No	Scientific name (Family)	Common names (Kurdish) English	Part used/ preparations	Traditional uses, to cure	Phytochemicals	Pharmacological effects
1	<i>Achillea ageratum</i> Linn. (Asteraceae)	(bejana zard) sweet maudlin	leaves, flowers	asthma, cough, diarrhea	fatty acids, phenolic acids, organic acids, phytosterols, flavonoids, terpenes including guaianolides (14, 15)	antibacterial, antioxidant, anti-inflammatory, cytotoxic effects (16)
2	<i>A. millefolium</i> Linn. (Asteraceae)	(bejan) yarrow	leaves, flowers / powder, boiling (Internal what means??)	bleeding, cold, ulcer, fever, to cure gynecological disorders	phytosterols, organic acids, phenolic acids, fatty acids, terpenes, including guaianolides, flavonoids (15, 17)	antispasmodic, hepatoprotective, (18)
3	<i>A. santolina</i> Linn. (Asteraceae)	(gulla beron)	Leaves, Flowers/ Fresh gum, Boiling (Internal)	gastrointestinal tract problems stomach diseases, ulcer	diterpenes, monoterpenes, sesquiterpenes, including sesquiterpene lactones, triterpenes, Flavonoids, Lignans, Essential oils (19)	antileishmanial, anticancer, antidiabetic (20)
4	<i>Alcea angulate</i> Freyn. (Malvaceae)	(gulle hiro) mallow	Flowers, Roots/ Boiling (internal)	Cold, Cough	Amino acids, Acidic polysaccharides, Flavonoids, Monosaccharides, pectinic acid, Syringic acid, p-Coumaric acid, Ferulic acid, p-Hydroxybenzoic acid, Caffeic acid, p-Hydroxyphenylacetic acid, Glucuronic acids, Rhamnose, and Galacturonic acid (21, 22)	hypolipidemic, cardioprotective, antimalarial (23)
5	<i>Allium ampeloprasum</i> Linn. (Amaryllidaceae)	(qurada) wild leek	Leaves, Bulbs/ Eaten	Antihypertensive	Irigenin, Sterebin A, Gerberinol, Rosmarinic acid, <i>ent</i> -epiafzelechin-(2 α -7,4 α -8)-catechin, Myristicanol B, Ramontoside, Urodienone, <i>trans</i> -Carvyl acetate, Epiafzelechin-(α -8)-pelargonidin-3'-glucoside, Boviquinone 4, Dihydrofukonolide, 2-(2-methylbutanoyl)-9-(3-methyl-2 <i>E</i> -pentenoyl)-2b,9a-dihydroxy4Z,10(14)-oplopadien-3-one, 6-Acetylfuranofukinol, Erinacine D, <i>N</i> - \square -glutamyl-S- <i>cis</i> -(1-propenyl) cysteine, <i>N</i> - \square -glutamyl-S-allyl cysteine, Terpenoids (24, 25)	
6	<i>Athaea officinalis</i> Linn. (Malvaceae)	(hero) marsh mallow	Flower/ Tea, Boiling	Cough	Alkaloids, Tannins, Triterpenes, Amino acids, Phenolic acids, Starch, Pectins, Asparagine, Coumarins, Phytosterols, Flavonoids, Mucilage (26-28)	Gastroprotective, antioxidant (29).
7	<i>Ammi visnaga</i> Linn. (Apiaceae)	(xalila) pick tooth	Seeds/ Boiling (Internal)	Vasodilator, Bronchodilator	Essential oils, Fatty acids, Flavonol, Flavonoids, Sterols, Phenols, Coumarins, γ -Pyrone, Visnagin and Khellin (30-34); Ammiol, Khellol, Khellinol, Visamminol, 4-Norvisamminol (35), Pimolin (III), 5,7-Dihydroxy-2-methyl- γ -pyrone-7- <i>O</i> -glucoside (36); <i>cis</i> -Khellactone-3'- β -D-glucopyranoside (37); Visnadin (38), Khellinin, Khellinone, Samidin, Dihydrosamidin, Visanginone (34); Ammoidin, Xanthotoxin, Psoralene, Bergapten (39, 40)	antioxidant, antifungal, antibacterial, and larvicidal (40).
8	<i>Anthemis des champs</i> (<i>Anthemis arvensis</i> Linn.) (Asteraceae)	(beibun) chamomile	Flowers/ (External, Internal)	Indigestion, Skin Whitening, Shortness of Breath, Diabetes	Essential oils, Polyphenolics, Glycosides, Quinic acid, Caffeic acid, Flavonoids including Quercetin and Apigenin-7-glucoside, α -Eudesmol, γ -Cadinene, γ -Cadinol, β -Pinene, Patuletin, Rosmarinic acid, Decanoic acid, Gentisic acid, Chlorogenic acid, 1,8-Cineole, T-Muurotol, α -Bisabolol, α -Bisabolene, Bisabolol oxide A,	antibacterial and anticancer (44)

9	<i>Artemisia absinthium</i> Mill. (Asteraceae)	(tolleke marane) wormwood absinthe	Flower/Decoction	Shortness of Breath, Diabetes	β -Farnesene, Chamazulene, α -Epicadinol, (+) Spathulenol, Germacrene D, Heptacosane, γ -Elemene, Caryophyllene oxide, Hexadecanoic acid ethyl ester, 9,12-Octadecadienoic acid (41-43)	Essential oil, Thujyl esters, α -Thujone (45)	
10	<i>Arum maculatum</i> Fisch. (Araceae)	(kardu) lord and ladies	Leaves/Boiling (Internal)	Gastrointestinal tract diseases	Phytol, Palmitic acid, Methyl linolenate, 9,12,15-Octadecatrienoate ester and Linolenate ester (37); Seventeen Chemicals have been isolated including Terpenes, Carbonyl Derivatives, Alcoholic compounds, Thiols, Fatty acid esters (38). More than 40 metabolites have been identified including aliphatic and aromatic compounds, mono-, sesquiterpenoids, nitrogen bearing compounds, and C5 branched chain components (39)		antioxidant, antimicrobial; toxic to humans (46)
11	<i>Asphodelus albus</i> Mill. (Asphodelaceae)	(astrelk) white asphodel	Roots/Powder	Anti-Dermatosis	Furan-3-ylmethyl ester and Essential oils (47).		
12	<i>Astragalus glaucacathus</i> Fisch. (Fabaceae)	(gueni) astragale	Leaves, Roots/Smoking (External)	Antiviral Infection	Amino acids, Flavonoids, Bioflavonoids, Saponins, Triterpene glycosides (31); Thidiazuron, Benzyladenine, Kinetin, Indole-3-acetic acid, Indole-3-butyric acid, Naphthalene acetic acid, Gibberellic acid, and 2,4-Dichlorophenoxyacetic acid (32); Carbohydrates and Hydrophilic biopolymers (33)	Mineral composition (Macro Micro elements), Vegetable oil, Tocopherols, Protein, Chlorogenic acids, Fatty acids, Amino acids (49, 50)	Anticancer, Antiviral, Antidiabetic, Anti-inflammatory, Hepatoprotective, Cardioprotective, Immunostimulant (48)
13	<i>Carlina acaulis</i> Linn. (Asteraceae)	(caw baza) stemless carline thistle	Flowers, Seeds/Powder, Juice	Liver Disease	Carotenoids, Anthocyanins, Polyphenols, Flavonoids, Tannins, Alkaloids (52, 53)		Anthelmintic, Antispasmodic, Diaphoretic, Diuretic, and Emmenagogic (51). against gastrointestinal-tract diseases, antioxidant, anti-inflammatory, antimicrobial, antidiarrheal, antiulcer, and laxative (54)
14	<i>Ceratonia siliqua</i> Linn. (Asteraceae)	(xrnuK) carob	Fruits/Boiling (Internal)	Abdominal Pain, Diarrhea	Phenolics, Flavonoids, Saponins, Tannins, Alkaloids, Protein, Glycosides, Carbohydrates, Ca ²⁺ , Mg ²⁺ , Mn ²⁺ , Cu ²⁺ , Na ⁺ , Fe ²⁺ , Zn ²⁺ , Se ²⁺ (55, 56)		Antimalarial, antibacterial, anthelmintic, antidiabetic (57)
15	<i>Cichorium intybus</i> Linn. (Asteraceae)	(ceq ceqe) chicory	Leaves, Bulbs, Flowers, Roots / Power, Boiling (Internal)	Liver, Gall Bladder	Polyphenolic compounds, Sterols, Esters, Saturated and Unsaturated fatty acids, Aromatic compounds (58)		antidiabetic Hepatoprotective, antioxidant, antitumor, anti-inflammatory (59)
16	<i>Cirsium vulgare</i> Petr. (Asteraceae)	(kenger) spear thistle	Leaves (Boiling (Internal))	gastrointestinal and kidney diseases	Flavonoids, Phenolic acids, Essential oils, Anthocyanins (60, 61)		antioxidant (61)
18	<i>Cota tinctoria</i> Linn. (Asteraceae)	(gulla hajila) yellow chamomile	Flowers (Tea, Boiling (Internal))	Fever, Inflammation	Tannins, Bioflavonoids, Phenolic compounds, Polysaccharides, Oligomeric procyanidins, Saponins, Vitamin C, Cardiotonic amines, Catecholamines, Ursolic acid, Purine (62, 63)		antidiarrheal, anti-inflammatory, immunomodulatory (62)
19	<i>Crataegus azarolus</i> Linn. (Rosaceae)	(gewij) hawthorn	Leaves, Fruits (Dried Parts as a tea, Fresh (Internal))	Heart diseases, Vasodilation	Polyphenolic and Phenolic acid derivatives, Flavonoids, Chicoric acid, Chlorogenic acid, Ketones, Terpenoids, Amides,		
20	<i>Crocus haussknechtii</i> Boiss. (Iridaceae)	(pishok) crocus	Flowers (Infusion (Internal))	Anti-Septic for gastritis and stomach problems			

21	<i>Dactylorhiza umbrosa</i> Kreutz. (Orchidaceae)	(salmka) marsh orchids or spotted orchids	Aerial Parts (Infusion (Internal))	Rheumatism	Flavones, carboxylic acids, Aldehydes (64) Phenolic and flavonoid derivatives, including quercetin, ascorbic acid, gallic acid (65-67)	
22	<i>Echinops viscosus</i> DC. (Asteraceae)	(kerteshi) globethistle	Fruits (Powder (Internal))	Git	Phenolic compounds (68)	Antioxidant, anticancer, antipyretic, hepatoprotective (69)
23	<i>Eryngium campestre</i> Linn. (Apiaceae)	(risheke) watling street thistle	Roots (Infusion (Internal))	Cough, Urinary problems, Infections, Kidney Pain	Essential oils, Terpenoids, Coumarins, Flavonoids, Acetylenes, Phenolic acids, Steroids, Saponins (70) Alkaloids (Berberine, Oxyberberine, Cryptopine, Fumarine, Parfumine, Protopine); Caffeic acid and Protocatechuic acid; Fatty acids, Volatile oils, Sesquiterpenoids, Aromatic hydrocarbons (71, 72)	Antibacterial, Antioxidant, Antitumor, Anti-inflammatory, and Hepatoprotective (73)
24	<i>Fumaria parviflora</i> Hook. (Papaveraceae)	(satere) fumitory	Leaves (Boiling (Internal))	Aerial Parts, Leaves	Terpenes, Glycosidic and Phenolic derivatives, Organic acids, Flavonoids (74) Several elements including Aluminum, Cobalt, Manganese, Iron, Nickel, Selenium, and Zinc (34); more than 50 chemical constituents have been extracted from leaves and analyzed by GC-MS (35); α -Amyrin, β -Amyrin, Cholesterol, Stigmasterol, β -Sitosterol, Ergosterol, β -Sitosterol-3-O-glucoside, Campesterol, Palmitic acid, Linoleic acid, Linolenic acid, Kampeferol-3-(6''-p-coumaroyl-O- β -D-glucoside), Chlorophyll A, Chlorophyll B, and Ethyl vanillin (36)	Antibacterial, Antioxidant, Antitumor, Anti-inflammatory, and Hepatoprotective (73)
25	<i>Lactuca serriola</i> Linn. (Asteraceae)	(talishk) dandelion root	Leaves (Raw (Internal))	Ulcer	Carvone, <i>cis</i> -dihydrocarvone, dihydrocarveol, limonene, 1,8-cineole, β -bourbonene, β -pinene, pulegone, piperitone, α -phellandrene, <i>trans</i> -caryophyllene, germacrene D (76); caffeic acid, Eriocitrin, Rosmarinic acid, Luteolin glucoside (77); Alkaloids, Flavonoids, Cardiac glycosides, Saponins, Steroids; more than 48 other constituents (78, 79); fat, proteins, carbohydrates, fiber, moisture, ash, Na ¹⁺ , Ca ²⁺ , K ¹⁺ , and Fe ²⁺ (80); Lignans (81)	
26	<i>Malva parviflora</i> Linn. (Malvaceae)	(tollaka) cheeseweed	Leaves (Boiling (Internal))	gastrointestinal trait diseases, Pain, Ulcer	phenolic acid and flavonoid derivatives, Pyrrol[de]phenanthridine, Amaryllidaceae alkaloid (lycorine), galanthamine (83, 84)	Anti-arthritic, anti-inflammatory (75)
27	<i>Mentha spicata</i> Linn. (Lamiaceae)	(pung) mint	Leaves (Boiling (Internal))	gastrointestinal trait diseases, carminative antispasmodic	Carotenoids, Phenols, Vitamin C, Phosphorus, Calcium (86-88)	Hepatoprotective, antiemetic, antianxiety (82)
28	<i>Narcissus tazetta</i> Linn. (Amaryllidaceae)	(nerges) paper white	Flower, Bulb (Boiling (Internal, External))	Headache, Cold, antiparasitic, Abortifacient	Essential oils and alkaloids such as Thebaine (Paramorphine), Morphine, Oripavine, Codeine, Rhoeadine, Papaverine; Phenolic compounds (90-92) Alkaloids (Morphine, Codeine, Rhoeadine, Thebaine, Papaverine), Phenolic compounds, Essential oils (90, 93, 94)	Anticancer, sedative, nervine tonic, anti-inflammatory (85)
29	<i>Nasturtium officinale</i> R.Br. (Brassicaceae)	(kuzalla) watercress	Leaves (Fresh (Internal))	Kidney problems	Alkaloids, Terpenoids, Flavonoids, Phenols, Saponins, Tannins, Cardioactive	Cardioprotective, Antibacterial, Anti-inflammatory, Anticancer, Antioxidant (89).
30	<i>Papaver bracteatum</i> Lindl. (Papaveraceae)	(gullalla sura) great scarlet poppy	Leaves and Flowers, (Raw (Internal))	appetite suppressant, Relaxant, Sedative		
31	<i>Papaver dubium</i> Linn. (Papaveraceae)	(gule shler) great scarlet poppy	Flowers and Leaves, Raw (Internal)	Sedative, relaxant for nervous System		Antimicrobial (95)
32	<i>Piperita mentha</i> Linn. (Lamiaceae)	(nah nah) peppermint	Leaves (Boiling (Internal))	gastrointestinal-trait, heart, brain, kidney diseases		Antimicrobial, antiviral, antioxidant antitumor (99)

33	<i>Plantago major</i> Linn. (Plantaginaceae)	(rkasha) rib grass	Leaves, Flowers, Latex (Powder, Extracts, hot infusion (Internal, External)	constipation, diarrhea, dysentery	glycosides, Carbohydrates (73, 96); Essential oils, Stilbenes, Phenolic lignans (97). Cardiac glycosides, Coumarins, Diterpenes, and Steroids (98)	anti-inflammatory, antitussive, anti- hemorrhagic, antiinfection, laxative, hemostatic, astringent, diuretic (102)
34	<i>Rheum ribes</i> Linn. (Polygonaceae)	(rewas) rhubarb	Roots, Seeds (Fresh, Powder (Internal)	diabetes, vitamin C Supplement, Cancer	Anthrones, Aloe-emodin, Emodin, Emodin glycoside, Physcion, Chrysophanol, Acylglucosides, Pyrones, Stilbenes, Flavonoids (103); Tannins and Phenolics (104); Alkaloids, Anthraquinones, Quercetin, Quinones (105); Palmitic acid, Hydrocarbons and long chain alkanes, including Heneicosane, Tricosane, Pentacosane, and Heptacosane (30)	Anti-inflammatory, anticancer, antibacterial, antiviral, expectorant (106)
35	<i>Salix alba</i> Linn. (Salicaceae)	(dar bi spi) white willow	Leaves (Decoction (Internal)	blood anticoagulant, anti- inflammatory, febrifuge	long-chain Alkanes, Alcohols, Carboxylic acids, Phenolics, Flavonoids, 3-O-Glycoside derivatives (107, 108)	
36	<i>Satureja khuzestanica</i> Jamzad (Lamiaceae)	(jatreh) summer	Stems, Leaves (Raw, Powder (Internal)	indigestion, headache, gastritis	Phenolic compounds (Carvacrol, Thymol), Essential oil, Triterpenes, Flavonoids, Glycosides, Steroids (109-111)	Antimicrobial, anticancer (112)
37	<i>Sinapis arvensis</i> Linn. (Brassicaceae)	(xertal)charlock mustard,field mustard	Stems, Flowers (Raw (Internal)	Laxative, Stomachache	Erythritol, 2-nitro-propanol, Cyclopentane Methyl Amine, 2- isopropylidene-N,N,5-trimethyl, Furfural, Thymol, Indole, Benzyl isothiocyanate, Cubenol, Essential oil, Monoterpenes, Sesquiterpenes, Sulfur- containing compounds (113, 114)	
38	<i>Smyrniium cordifolium</i> Boiss (Araceae)	horse parsley (qalandor)	Stem (Raw (Internal)	Laxative, Stomachache, Headache	Flavonoids, Monoterpenes, Essential oil (115)	
39	<i>Taraxacum officinale</i> Koch. (Asteraceae)	(talishka) dandelion root	Leaves (Decoction (Internal)	Tuberculosis	Alkaloids, Terpenoids, Phenols, Flavonoids, Glycosides, Saponins, Tannins (116, 117)	Anticancer, Antioxidant, Anti- inflammatory, Anti- hyperglycemic, Anticoagulant, Analgesic, Choleric, Diuretic (118).
40	<i>Thymus vulgaris</i> Linn. (Lamiaceae)	(catre) thyme	Leaves, Flowers (Boiling (Internal)	Gastrointestinal - trait, heart, and kidney diseases, inflammation, diabetes, obesity	Essential oil (Thymol, p- Cymene, Terpinene, Linalool, Carvacrol), 4-Hydroxybenzoic acid, Vanillic acid, Catechin, Epicatechin, Syringic acid, Gallic acid, Chlorogenic acid, Protocatechuic acid, Caffeic acid, Catechin, trans-Cinnamic acid, γ -Coumaric acid, trans- Ferulic acid, Kaempferol, Myricetin, Quercetin, Tocopherol, α -Tocopherol, β - Tocopherol, γ -Tocopherol, δ - Tocopherol, Campesterol, β - Sitosterol, Ergosterol, and Stigmasterol (119, 120).	Cytotoxic, antioxidant, antimicrobial, mosquitocidal (121).

The structures of the most important Kodo Mountain are displayed in the following Figs 8-11.




Fig. 8: Chemical structure of some isolated chemicals from *Mentha spicata* L. (76, 77).




Fig. 9: Structure of the main isolated constituents in *Malvaceae* genus (21, 22, 26-28, 36).



Fig 10. Chemical structure of some flavonoids isolated from *Anthemis des champs*(41-43)

Fig. 11: Structures of the main alkaloids found in different species of plants in Kodo mountain (90-94).

Conclusion

Our first ethnopharmacological review of medicinal plants used in the Kodo Mountain, which was based on interviews with local healers, demonstrated that traditional medicine is still widely practiced by this group of Kurds, as well as by other communities of Middle East countries. This fund of knowledge has been passed down through generations only orally; therefore, it is important to keep it in a well-organized form in written documentation, how does this paper want to be. The effectiveness of medicinal plants growing in the Kodo mountain for a wide range of human ailments is not only supported but their prolonged use over time but also by the scientific literature reporting the biological properties of isolated bioactive compounds. However, most data have been obtained from samples of plants collected in other regions; therefore, given the frequent dependence of the contents of secondary metabolites on the plant geographical origin, it is recommendable to examine species native to the Kodo Mountain. Moreover, before considering the local herbal remedies as potential targets for the research of new drugs, more investigations are needed, such as the exploration of the mechanisms of action, and in vivo studies aimed at defining the pharmacokinetic profile, safety, toxic and side effects.

In summary, we believe that this paper demonstrates the great potential of the medicinal plants growing in the Kodo mountain as sources of products for different purposes and applications.

Conflict of Interest

All the authors declare that there are no conflicts of interest.

Acknowledgement

The authors are grateful to Tishk International University (TIU) for supporting this investigation. We also greatly appreciate the Kurdish informants' traditional knowledge of the therapeutic applications of the plants growing in Kodo Mountain.

References

1. J. W. Harshberger, The purposes of ethnobotany. *Bot. Gaz.* **21**, 146-154 (1896).
2. F. O. Abdullah, F. H. Hussain, M. Clericuzio, A. Porta, G. Vidari, A new iridoid dimer and other constituents from the traditional Kurdish plant *Pterocephalus nestorianus* Nábělek. *Chem. Biodivers.* **14**, e1600281 (2017).
3. F. O. Abdullah, F. H. Hussain, L. I. Cucca, G. Vidari, Phytochemical investigation and antioxidant effects of different solvent extracts of *Pterocephalus nestorianus* Nab. growing in Kurdistan Region-Iraq. *Science Journal of University of Zakho* **6**, 21-25 (2018).
4. F. O. Abdullah *et al.*, Composition, antifungal and antiproliferative activities of the hydrodistilled oils from leaves and flower heads of *Pterocephalus nestorianus* Nábělek. *Chem. Biodivers.* **14**, e1700009 (2017).
5. F. O. Abdullah *et al.*, Iridoids Isolation from a Phytochemical Study of the Medicinal Plant *Teucrium parviflorum* Collected in Iraqi Kurdistan. *Molecules* **27**, 5963 (2022).
6. H. I. M. Amin *et al.*, Chemical composition and antifungal activity of essential oils from flowers, leaves, rhizomes, and bulbs of the wild Iraqi Kurdish plant *Iris persica*. *Nat. Prod. Commun.* **12**, 1934578X1701200334 (2017).
7. H. I. M. Amin, F. H. Hussain, M. Maggolini, G. Vidari, Bioactive constituents from the traditional Kurdish plant *Iris persica*. *Nat. Prod. Commun.* **13**, 1934578X1801300907 (2018).
8. H. I. M. Amin *et al.*, Phytochemistry and biological activities of *Iris* species growing in Iraqi Kurdistan and phenolic constituents of the traditional plant *Iris postii*. *Molecules* **26**, 264 (2021).
9. M. F. Ibrahim, F. H. S. Hussain, G. Zanoni, G. Vidari, The main constituents of *Tulipa systola* Stapf. roots and flowers; their antioxidant activities. *Nat. Prod. Res.* **31**, 2001-2007 (2017).
10. H. I. M. Amin *et al.*, Phytochemistry of *Verbascum* Species Growing in Iraqi Kurdistan and Bioactive Iridoids from the Flowers of *Verbascum calvum*. *Plants* **9**, 1066 (2020).
11. P. Prakash *et al.*, Documentation of commonly used ethnoveterinary medicines from wild plants of the high mountains in Shimla District, Himachal Pradesh, India. *Horticulturae* **7**, 351 (2021).
12. P. Prakash *et al.*, Therapeutic uses of wild plants by rural inhabitants of Maraog region in district Shimla, Himachal Pradesh, India. *Horticulturae* **7**, 343 (2021).
13. A. A. Othman, R. Gloaguen, Automatic extraction and size distribution of landslides in Kurdistan Region, NE Iraq. *Remote Sensing* **5**, 2389-2410 (2013).
14. L. Tebaa *et al.*, Algicidal effects of *Achillea ageratum* L. and *Origanum compactum* Benth. plant extracts on growth of *Microcystis aeruginosa*. *Appl Ecol Environ Res* **15**, 719-728 (2017).

15. M. Strzypek-Gomólka, K. Gawel-Bęben, W. Kukula-Koch, Achillea species as sources of active phytochemicals for dermatological and cosmetic applications. *Oxidative medicine and cellular longevity* **2021**, (2021).
16. L. El Bouzidi *et al.*, Essential oil composition and antimicrobial activity of wild and cultivated Moroccan Achillea ageratium L.: a rare and threatened medicinal species. *Chem. Biodivers.* **9**, 598-605 (2012).
17. M. Farajpour, M. Ebrahimi, A. Baghizadeh, M. Aalifar, Phytochemical and yield variation among Iranian Achillea millefolium accessions. *HortScience* **52**, 827-830 (2017).
18. S. I. Ali, B. Gopalakrishnan, V. Venkatesalu, Pharmacognosy, phytochemistry and pharmacological properties of Achillea millefolium L.: a review. *Phytother. Res.* **31**, 1140-1161 (2017).
19. S. Kumar, R. Kumari, Pharmacological Activities of Achillea santolina: A Review. *Am J Pharmacol Pharmacother* **8**, 13 (2021).
20. F. Ayrom, S. Rasouli, B. Shemshadi, In vitro antileishmanial activity of Achillea santolina essential oil against Leishmania infantum promastigote by methylthiazole tetrazolium (MTT) and trypan blue colorimetric methods. *Archives of Razi Institute* **76**, 529 (2021).
21. Z. Fahimi, S. Soltanbeigi, K. Asadollahi, M. Kaffashi, M. Kaffashian, Effects of Alcoholic Extract of Stem and Aerial Parts of the Alcea angulate Plant on Aorta Atherosclerotic Plaque Formation in Rabbit. *Archives of Medicine* **10**, 0-0 (2018).
22. P. Saha, Brac University, (2019).
23. Z. Fahimi, J. Cheraghi, A. Pilehvarian, K. Sayehmiri, A. Khosravi, Effects of Alcea angulata root alcoholic extract on blood lipid of male rabbit. (2012).
24. P. García-Herrera *et al.*, Nutrients, phytochemicals and antioxidant activity in wild populations of Allium ampeloprasum L., a valuable underutilized vegetable. *Food Res. Int.* **62**, 272-279 (2014).
25. C. Ceccanti *et al.*, Comparative phytochemical profile of the elephant garlic (Allium ampeloprasum var. holmense) and the common garlic (Allium sativum) from the Val di Chiana area (Tuscany, Italy) before and after in vitro gastrointestinal digestion. *Food Chem.* **338**, 128011 (2021).
26. Y. J. Park, N. S. Kim, R. Sathasivam, Y. S. Chung, S. U. Park, Impact of copper treatment on phenylpropanoid biosynthesis in adventitious root culture of Althaea officinalis L. *Prep Biochem Biotechnol* **52**, 283-291 (2022).
27. A. Pirmohammadi, A. Abdali Mashhadi, A. Koochekzadeh, A. Lotf Jalal Abadi, A. Siahpoush, Effect of osmopriming and zinc foliar spraying on yield components and mucilage of flower in Marshmallow (Althaea officinalis L.) medicinal plant. *Iranian Journal of Horticultural Science* **52**, 877-887 (2022).
28. M. Husain, A. Wadud, G. Sofi, S. Perveen, K. A. Hafeez, Physicochemical standardization of mucilage obtained from Althaea officinalis Linn-Root. *Pharmacognosy Magazine* **15**, 155 (2019).
29. M. Mahboubi, Marsh mallow (Althaea officinalis L.) and its potency in the treatment of cough. *Complementary medicine research* **27**, 174-183 (2020).
30. M.-R. Nikbakht, S. Esnaashari, F. Heshmati Afshar, Chemical composition and general toxicity of essential oil extracted from the stalks and flowers of Rheum ribes L. growing in Iran. *Journal of Reports in Pharmaceutical Sciences* **2**, 165-170 (2013).
31. H. Kızıltaş *et al.*, LC-HRMS profiling of phytochemicals, antidiabetic, anticholinergic and antioxidant activities of evaporated ethanol extract of Astragalus brachycalyx Fischer. *Journal of Chemical Metrology*, (2021).
32. A. Yildirim, E. Uyar, A. Turker, In vitro culture of endemic Astragalus gymnolobus Fischer and comparison of its antibacterial, antioxidant, and phenolic profiles with field grown plants. *Journal of Agricultural Science and Technology* **22**, 815-828 (2020).
33. E. Ahmadi, H. Rezadoost, M. M. Farimani, Isolation, characterization, and antioxidant activity of neutral carbohydrates from Astragalus arbusculus gum. *S. Afr. J. Bot.* **146**, 669-675 (2022).
34. B. A. Sadee, Determination of Elements in Indigenous Vegetables Using ICP-MS. *Cihan University-Erbil Scientific Journal* **6**, 26-31 (2022).
35. E. A. Khoris, E. M. El-Sherbeny, trying to use antibiotics alternatives to raise immune efficiency and growth performance in tilapia nilotica. *Assiut Vet Med J* **68**, 44 (2022).
36. A. Abdel-Ghani, H. Hassan, A. M. Elshazly, Phytochemical and biological study of Malva parviflora L. grown in Egypt. *Zagazig Journal of Pharmaceutical Sciences* **22**, 17 (2013).
37. H. S. Al-Shmgani, Z. H. M. Kadri, M. M. Al-Halbosi, Y. H. Dewir, Phytochemical analysis, cytotoxicity and antioxidant activity of cuckoo pint (Arum maculatum) leaf extract. *Acta Biol Szeged* **63**, 119 (2019).

38. V. Kochmarov *et al.*, Exploration of collagenase, cyclooxygenases, angiogenesis and free radical processes as the putative pharmacological targets of *Arum maculatum* L. *Biotechnology & Biotechnological Equipment* **34**, 126 (2020).
39. G. Marotz-Clausen *et al.*, Incomplete synchrony of inflorescence scent and temperature patterns in *Arum maculatum* L.(Araceae). *Phytochemistry* **154**, 77 (2018).
40. N. Khalil, M. Bishr, S. Desouky, O. Salama, *Ammi visnaga* L., a potential medicinal plant: A review. *Molecules* **25**, 301 (2020).
41. I. H. Mohammed, A. T. Hameed, H. F. Salman, Phytochemical and Biological of *Anthemis nobilis* (Asteraceae family) a Native Herbs of Iraq. *Sys Rev Pharm* **11**, 458-461 (2020).
42. A. Raal *et al.*, The Phytochemical Profile and Anticancer Activity of *Anthemis tinctoria* and *Angelica sylvestris* Used in Estonian Ethnomedicine. *Plants* **11**, 994 (2022).
43. S. Adhikari, I. C. Burke, S. D. Eigenbrode, Mayweed chamomile (*Anthemis cotula* L.) biology and management—a review of an emerging global invader. *Weed Res.* **60**, 313-322 (2020).
44. L. Riccobono, A. Maggio, M. Bruno, V. Spadaro, F. M. Raimondo, Chemical composition and antimicrobial activity of the essential oils of some species of *Anthemis* sect. *Anthemis* (Asteraceae) from Sicily. *Nat. Prod. Res.* **31**, 2759-2767 (2017).
45. A. Szopa *et al.*, *Artemisia absinthium* L.—Importance in the history of medicine, the latest advances in phytochemistry and therapeutical, cosmetological and culinary uses. *Plants* **9**, 1063 (2020).
46. R. Farahmandfar, R. Esmaeilzadeh Kenari, M. Asnaashari, D. Shahrapour, T. Bakhshandeh, Bioactive compounds, antioxidant and antimicrobial activities of *Arum maculatum* leaves extracts as affected by various solvents and extraction methods. *Food science & nutrition* **7**, 465-475 (2019).
47. M. S. Dekić, E. S. Selimović, New Natural Products from *Asphodelus albus* MILL. Essential Oil. *Chem. Biodivers.* **18**, e2100103 (2021).
48. X. Li *et al.*, A review of recent research progress on the *astragalus* genus. *Molecules* **19**, 18850-18880 (2014).
49. M. Strzemeski *et al.*, Nitrogen fertilisation decreases the yield of bioactive compounds in *Carlina acaulis* L. grown in the field. *Industrial Crops and Products* **170**, 113698 (2021).
50. M. Strzemeski *et al.*, Morphological, Anatomical, and Phytochemical Studies of *Carlina acaulis* L. *Cypselia. Int. J. Mol. Sci.* **21**, (2020).
51. M. Strzemeski, M. Wójciak-Kosior, I. Sowa, D. Załuski, R. Verpoorte, Historical and traditional medical applications of *Carlina acaulis* L.-A critical ethnopharmacological review. *J. Ethnopharmacol.* **239**, 111842 (2019).
52. K. Ben Othmen, J. M. Garcia-Beltrán, W. Elfalleh, M. Haddad, M. Á. Esteban, Phytochemical compounds and biological properties of carob pods (*Ceratonia siliqua* L.) extracts at different ripening stages. *Waste and Biomass Valorization* **12**, 4975-4990 (2021).
53. I. Lakkab *et al.*, *Ceratonia siliqua* L. seed peels: Phytochemical profile, antioxidant activity, and effect on mood disorders. *J. Funct. Foods* **54**, 457-465 (2019).
54. K. Rūbi *et al.*, Chemical constituents and pharmacological actions of carob pods and leaves (*Ceratonia siliqua* L.) on the gastrointestinal tract: A review. *Biomed. Pharmacother.* **93**, 522-528 (2017).
55. Z. K. Abbas *et al.*, Phytochemical, antioxidant and mineral composition of hydroalcoholic extract of chicory (*Cichorium intybus* L.) leaves. *Saudi J. Biol. Sci.* **22**, 322-326 (2015).
56. M. Arya, B. R. Singh, G. Taj, Phytochemical screening and quantitative analysis of *Cichorium intybus* L.(Chicory) plants from region of Uttarakhand. *The Pharma Innovation Journal* **11**, 230-235 (2022).
57. R. Street, J. Sidana, G. Prinsloo, *Cichorium intybus*: traditional uses, phytochemistry, pharmacology and toxicology evidence-based. *J. Complement. Altern. Med.* **15**, (2013).
58. S. Kadan *et al.*, *Gundelia tournefortii*: Fractionation, Chemical Composition and GLUT4 Translocation Enhancement in Muscle Cell Line. *Molecules* **26**, 3785 (2021).
59. W. Luo *et al.*, Recent research progress of *Cirsium* medicinal plants in China. *J. Ethnopharmacol.* **280**, 114475 (2021).
60. S. Shamloo, S. J. Marandi, G. Tajadod, A. Majd, R. Rahimi, Cytotoxic effect of hydroalcoholic extract of *Cota tinctoria* (L.) J. Gay on AGS and Hep-G2 cancer cell lines. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas* **21**, (2022).
61. M. B. Bahadori *et al.*, Determination of phenolics composition, antioxidant activity, and therapeutic potential of Golden marguerite (*Cota tinctoria*). *Journal of Food Measurement and Characterization* **15**, 3314 (2021).
62. E. Abu-Gharbieh, N. G. Shehab, Therapeutic potentials of *Crataegus azarolus* var. *eu-azarolus*

- Maire leaves and its isolated compounds. *BMC complementary and alternative medicine* **17**, 1 (2017).
63. H. H. Balaky *et al.*, Estimation of total tannin and total phenolic content in plant (*Crataegus azarolus* L) by orbital shaker technique. *International Journal of Agriculture Environment and Food Sciences* **5**, 1 (2021).
64. N. Zivyar, G. Bagherzade, M. Moudi, M. Manzari Tavakoli, Evaluation of the green synthesis, characterization and antibacterial activity of silver nanoparticles from corm extract of *Crocus sativus* var. *Hausknechtii*. *Journal of Horticulture and Postharvest Research* **4**, 19-32 (2021).
65. A. Khuroo, G. Shapoo, Z. Kaloo, A. Ganie, S. Singh, *Dactylorhiza umbrosa* (Kar. & Kir.) Nevski (Orchidaceae): an addition to flora of India from Kashmir Himalaya. *Check List* **12**, 1-5 (2016).
66. H. KIZILTAŞ, E. Suat, D. YILDIZ, S. M. PINAR, Evaluation of Antioxidant Properties, Trace Element and Mineral Composition of *Dactylorhiza umbrosa* (Kar. & Kir.) Nevski (Orchidaceae). *Journal of the Institute of Science and Technology* **9**, 2148-2156 (2019).
67. L. V. Averyanov, A review of the genus *Dactylorhiza*. *Orchid biology. Reviews and perspectives* **5**, 159-206 (1990).
68. H. M. Abdallah, H. Z. Asfour, A. M. El-Halawany, M. A. Elfaky, Saudi plants as a source of potential β -lactamase inhibitors. *Pak. J. Pharm. Sci.* **31**, (2018).
69. H. Bitew, A. Hymete, The genus *Echinops*: Phytochemistry and biological activities: A review. *Front. Pharmacol.* **10**, 1234 (2019).
70. B. Soumia, *Eryngium campestre* L.: Polyphenolic and flavonoid compounds; Applications to health and disease. *Polyphenols: Mechanisms of Action in Human Health and Disease*, 69-79 (2018).
71. A. Bhargava, P. Shrivastava, A. Tilwari, HPTLC analysis of *Fumaria parviflora* (Lam.) methanolic extract of whole plant. *Future Journal of Pharmaceutical Sciences* **7**, 1-9 (2021).
72. S. Kumar, A. K. Sharma, A. Kamboj, *Fumaria parviflora* Lam.(Fumitory): A traditional herbal medicine with modern evidence. *Asian Journal of Pharmacy and Pharmacology* **3**, 200-207 (2017).
73. R. Zhang, Q. Guo, E. J. Kennelly, C. Long, X. Chai, Diverse alkaloids and biological activities of *Fumaria* (Papaveraceae): An ethnomedicinal group. *Fitoterapia* **146**, 104697 (2020).
74. N. H. A. Fatah, Y. Amen, F. Abdel Bar, A. F. Halim, H.-E. A. Saad, Antioxidants and α -glucosidase Inhibitors from *Lactuca serriola* L. *Rec. Nat. Prod* **14**, 410-415 (2020).
75. G. B. Martínez-Hernández *et al.*, Anti-arthritis and anti-inflammatory effects of extract and fractions of *Malva parviflora* in a mono-arthritis model induced with kaolin/carrageenan. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **393**, 1281-1291 (2020).
76. M. Mahboubi, *Mentha spicata* L. essential oil, phytochemistry and its effectiveness in flatulence. *Journal of Traditional and Complementary Medicine* **11**, 75-81 (2021).
77. S. R. Kanatt, R. Chander, A. Sharma, Antioxidant potential of mint (*Mentha spicata* L.) in radiation-processed lamb meat. *Food Chem.* **100**, 451-458 (2007).
78. S. Zaidi, P. Dahiya, In vitro antimicrobial activity, phytochemical analysis and total phenolic content of essential oil from *Mentha spicata* and *Mentha piperita*. *International Food Research Journal* **22**, 2440 (2015).
79. P. Jain, A. Soni, P. Jain, J. Bhawsar, Phytochemical analysis of *Mentha spicata* plant extract using UV-VIS, FTIR and GC/MS technique. *J. Chem. Pharm. Res.* **8**, 1-6 (2016).
80. A. M. E. Sulieman, S. E. Abdelrahman, A. Abdel Rahim, Phytochemical analysis of local spearmint (*Mentha spicata*) leaves and detection of the antimicrobial activity of its oil. *Journal of Microbiology Research* **1**, 1-4 (2011).
81. G. Mahendran, S. K. Verma, L.-U. Rahman, The traditional uses, phytochemistry and pharmacology of spearmint (*Mentha spicata* L.): A review. *J. Ethnopharmacol.* **278**, 114266 (2021).
82. J. Ahamad, Aroma Profile and α -Glucosidase Inhibitory Activity of Essential Oil of *Mentha spicata* Leaves. *Journal of Essential Oil Bearing Plants* **24**, 1042-1048 (2021).
83. Ç. Karakoyun *et al.*, A brief up-to-date overview of Amaryllidaceae alkaloids: Phytochemical studies of *Narcissus tazetta* subsp. *tazetta* L., collected in Turkey. *Nat. Prod. Commun.* **14**, 1934578X19872906 (2019).
84. A. Tarakemeh *et al.*, Screening of Amaryllidaceae alkaloids in bulbs and tissue cultures of *Narcissus papyraceus* and four varieties of *N. tazetta*. *J. Pharm. Biomed. Anal.* **172**, 230-237 (2019).
85. W. H. Talib, A. M. Mahasneh, Antimicrobial, Cytotoxicity and Phytochemical Screening of Jordanian Plants Used in Traditional Medicine. *Molecules* **15**, 1811-1824 (2010).

86. S. Zaman *et al.*, Determination of genetic diversity, sinigrin contents, and elicitors-induced enhancement of sinigrin in *Nasturtium officinale* L. *Turkish Journal of Agriculture and Forestry* **46**, 59-72 (2022).
87. S. Kyriakou *et al.*, Evaluation of Bioactive Properties of Lipophilic Fractions of Edible and Non-Edible Parts of *Nasturtium officinale* (Watercress) in a Model of Human Malignant Melanoma Cells. *Pharmaceuticals* **15**, 141 (2022).
88. G. Pignata *et al.*, Understanding the Postharvest Phytochemical Composition Fates of Packaged Watercress (*Nasturtium officinale* R. Br.) Grown in a Floating System and Treated with *Bacillus subtilis* as PGPR. *Plants* **11**, 589 (2022).
89. M. Klimek-Szczykutowicz, A. Szopa, H. Ekiert, Chemical composition, traditional and professional use in medicine, application in environmental protection, position in food and cosmetics industries, and biotechnological studies of *Nasturtium officinale* (watercress)—a review. *Fitoterapia* **129**, 283-292 (2018).
90. M. Butnariu *et al.*, Papaver Plants: Current Insights on Phytochemical and Nutritional Composition Along with Biotechnological Applications. *Oxidative Medicine and Cellular Longevity* **2022**, (2022).
91. A. Qaderi *et al.*, Molecular diversity and phytochemical variability in the Iranian poppy (*Papaver bracteatum* Lindl.): A baseline for conservation and utilization in future breeding programmes. *Industrial Crops and Products* **130**, 237-247 (2019).
92. A. Habibzade, N. Moshtaghi, S. Malekzadeh-Shafaroudi, Phytochemical investigation among different populations of *Papaver bracteatum* Lindl. using ultrasonic-assisted extraction method followed by HPLC analysis. *Journal of Medicinal Plants* **20**, 23-33 (2021).
93. A. Mat *et al.*, Alkaloids and bioactivity of *Papaver dubium* subsp. *dubium* and *P. dubium* subsp. *laevigatum*. *Nat. Prod. Lett.* **14**, 205-210 (2000).
94. H. Da Cheng, G. Xiao-Jie, G. Pei, Phytochemical and biological research of *Papaver* pharmaceutical resources. *Medicinal Plants: Chemistry Biology and Omics*, 217-251 (2015).
95. P. Razaghi, D. Zafari, Characterization of fungi causing lesion blight on *Papaver dubium* in Iran. *Antonie Van Leeuwenhoek* **111**, 437-455 (2018).
96. A. N. Adham, Comparative extraction methods, phytochemical constituents, fluorescence analysis and HPLC validation of rosmarinic acid content in *Mentha Piperita*, *Mentha longifolia* and *Osimum basilicum*. *Natural Products: An Indian Journal* **11**, (2015).
97. G. Mahendran, L.-U. Rahman, Ethnomedicinal, phytochemical and pharmacological updates on Peppermint (*Mentha × piperita* L.)—A review. *Phytother. Res.* **34**, 2088-2139 (2020).
98. S. R. Patil, R. S. Patil, A. G. Godghate, *Mentha Piperita* Linn: Phytochemical, Antibacterial And Dipterian Adulticidal Approach. *International Journal of Pharmacy and Pharmaceutical Sciences* **8**, 352-355 (2016).
99. D. L. McKay, J. B. Blumberg, A review of the bioactivity and potential health benefits of peppermint tea (*Mentha piperita* L.). *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives* **20**, 619-633 (2006).
100. Y. Najafian, S. S. Hamed, M. K. Farshchi, Z. Feyzabadi, *Plantago major* in Traditional Persian Medicine and modern phytotherapy: a narrative review. *Electron Physician* **10**, 6390-6399 (2018).
101. M. A. Soliman, T. M. Galal, M. A. Naeim, A. A. Khalafallah, Seasonal Variation in the Secondary Metabolites and Antimicrobial Activity of *Plantago major* L. from Egyptian Heterogenic Habitats. *Egypt J Bot* **62**, 255-273 (2022).
102. Y. Najafian, S. S. Hamed, M. K. Farshchi, Z. Feyzabadi, *Plantago major* in Traditional Persian Medicine and modern phytotherapy: a narrative review. *Electronic physician* **10**, 6390 (2018).
103. P. Singh, M. Rawat, Phytochemistry and biological activity perspectives of *Rheum* species. *The natural products journal* **6**, 84-93 (2016).
104. T. TAŞKIN, G. Bulut, Qualitative and quantitative phytochemical analysis and in-vitro biological activity of *Rheum ribes* L. different parts. *Istanbul Journal of Pharmacy* **49**, 7-13 (2019).
105. A. N. Adham, A. M. Naqishbandi, HPLC analysis and antidiabetic effect of *Rheum ribes* root in type 2 diabetic patients. *Zanco Journal of Medical Sciences (Zanco J Med Sci)* **19**, 957-964 (2015).
106. Z. Keshavarzi *et al.*, A Review on the Phytochemistry, Pharmacology, and Therapeutic Effects of *Rheum ribes*. *Natural Products and Human Diseases*, 447-461 (2021).
107. A. L. Santos, M. G. Soares, L. S. de Medeiros, M. J. P. Ferreira, P. Sartorelli, Identification of flavonoid-3-O-glycosides from leaves of *Casearia arborea* (Salicaceae) by UHPLC-DAD-ESI-HRMS/MS combined with molecular

- networking and NMR. *Phytochem. Anal.* **32**, 891-898 (2021).
108. E. Autor *et al.*, Extraction of Phenolic Compounds from Populus Salicaceae Bark. *Biomolecules* **12**, 539 (2022).
109. N. Moradi-Maram, D. Dastan, A. Nili-Ahmadabadi, Protective Effect of Various Extracts of Allium hirtifolium and Satureja khuzestanica Plants on AAPH-Induced Oxidative Hemolysis. *Avicenna Journal of Pharmaceutical Research* **2**, 1-7 (2021).
110. B. Sadeghi-Nejad, A. Rezaei-Matehkolaei, S. Y. Naanaie, Isolation and antifungal activity evaluation of Satureja khuzestanica Jamzad extract against some clinically important dermatophytes. *J. Mycol. Med.* **27**, 554-560 (2017).
111. F. Sefidkon, Z. Emami Bistgani, Integrative review on ethnobotany, essential oil, phytochemical, agronomy, molecular and pharmacological properties of Satureja species. *J. Essent. Oil Res.* **33**, 114-132 (2021).
112. A. Khaledi, M. Meskini, A systematic review of the effects of Satureja khuzestanica Jamzad and Zataria multiflora Boiss against Pseudomonas aeruginosa. *Iran J Med Sci* **45**, 83 (2020).
113. A. F. Al-Rubaye, M. J. Kadhim, I. H. Hameed, Determination of bioactive chemical composition of methanolic leaves extract of Sinapis arvensis using GC-MS technique. *Int J Toxicol Pharmacol Res* **9**, 163-178 (2017).
114. J. S. Rad, M. H. Alfatemi, M. S. Rad, D. Sen, Phytochemical and antimicrobial evaluation of the essential oils and antioxidant activity of aqueous extracts from flower and stem of Sinapis arvensis L. *American Journal of Advanced Drug Delivery* **1**, 001-010 (2013).
115. N. Abbasi *et al.*, Protective effects of smyrnium cordifolium boiss essential oil on pentylenetetrazol-induced seizures in mice: Involvement of benzodiazepine and opioid antagonists. *J. Biol. Regul. Homeost. Agents* **31**, 683-689 (2017).
116. M. A. Mir, S. Sawhney, M. Jassal, Qualitative and quantitative analysis of phytochemicals of Taraxacum officinale. *Wudpecker Journal of Pharmacy and Pharmacology* **2**, 001-005 (2013).
117. D. Jedrejek, B. Lis, A. Rolnik, A. Stochmal, B. Olas, Comparative phytochemical, cytotoxicity, antioxidant and haemostatic studies of Taraxacum officinale root preparations. *Food Chem. Toxicol.* **126**, 233-247 (2019).
118. K. Schütz, R. Carle, A. Schieber, Taraxacum—a review on its phytochemical and pharmacological profile. *J. Ethnopharmacol.* **107**, 313-323 (2006).
119. P. Daga, S. R. Vaishnav, A. Dalmia, A. W. Tumaney, Extraction, fatty acid profile, phytochemical composition and antioxidant activities of fixed oils from spices belonging to Apiaceae and Lamiaceae family. *J. Food Sci. Technol.* **59**, 518-531 (2022).
120. R. Tardugno *et al.*, Thymus vulgaris L. essential oils from Emilia Romagna Apennines (Italy): phytochemical composition and antimicrobial activity on food-borne pathogens. *Nat Prod Res* **36**, 837-842 (2022).
121. S. Sertel, T. Eichhorn, P. K. Plinkert, T. Efferth, Cytotoxicity of Thymus vulgaris essential oil towards human oral cavity squamous cell carcinoma. *Anticancer Res.* **31**, 81-87 (2011).